

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 0 862 566 B1**

(12)

# EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention  
of the grant of the patent:  
**12.01.2000 Bulletin 2000/02**

(21) Application number: **96937248.1**

(22) Date of filing: **25.10.1996**

(51) Int Cl.7: **C07D 401/04, C07D 405/14,  
C07D 401/14, C07D 413/14,  
C07D 471/14, C07D 417/14,  
C07D 409/14, C07D 471/04**

(86) International application number:  
**PCT/EP96/04660**

(87) International publication number:  
**WO 97/16440 (09.05.1997 Gazette 1997/20)**

(54) **1-(1,2-DISUBSTITUTED PIPERIDINYL)-4-SUBSTITUTED PIPERAZINE DERIVATIVES**

**1-(1,2-DISUBSTITUIERTE PIPERIDINYL)-4-SUBSTITUIERTE PIPERAZIN DERIVATE**

**DERIVES DE PIPERAZINE A SUBSTITUTION EN POSITIONS 1-(DIPERIDINYLE A  
DISUBSTITUTION EN POSITIONS 1,2)-4**

(84) Designated Contracting States:  
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU NL  
PT SE**  
Designated Extension States:  
**LT LV SI**

(30) Priority: **30.10.1995 EP 95202929**

(43) Date of publication of application:  
**09.09.1998 Bulletin 1998/37**

(73) Proprietor: **JANSSEN PHARMACEUTICA N.V.  
2340 Beerse (BE)**

(72) Inventors:  
• **JANSSENS, Frans, Eduard  
B-2820 Bonheiden (BE)**  
• **SOMMEN, François, Maria  
B-2323 Wortel (BE)**

• **SURLERAUX, Dominique, Louis, Nestor,  
Ghislaine  
B-1830 Machelen (BE)**  
• **LEENAERTS, Joseph, Elisabeth  
B-2310 Rijkevorsel (BE)**  
• **VAN ROOSBROECK, Yves, Emiel, Maria  
B-2220 Heist-op den-Berg (BE)**

(74) Representative: **Quaghebeur, Luc  
Janssen Pharmaceutica N.V.,  
Patent Department,  
Turnhoutseweg 30  
2340 Beerse (BE)**

(56) References cited:  
**EP-A- 0 512 901 EP-A- 0 532 456  
EP-A- 0 625 509 EP-A- 0 655 442  
WO-A-95/11895**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 0 862 566 B1**

## Description


[0001] This invention concerns 1-(1,2-disubstituted piperidiny)-4-substituted piperazine derivatives having tachykinin antagonistic activity, in particular substance P antagonistic activity, and their preparation; it further relates to compositions comprising them, as well as their use as a medicine.

[0002] Substance P is a naturally occurring neuropeptide of the tachykinin family. There are ample studies showing that substance P and other tachykinins are involved in a variety of biological actions, and therefore, play an essential role in various disorders (Regoli et al., Pharmacological Reviews 46(4), 1994, p. 551-599, "Receptors and Antagonists for Substance P and Related Peptides"). The development of tachykinin antagonists has led to date to a series of peptide compounds of which might be anticipated that they are metabolically too labile to be employed as pharmaceutically active substances (Longmore J. et al., DN&P 8(1), February 1995, p. 5-23, "Neurokinin Receptors"). The present invention concerns nonpeptide tachykinin antagonists, in particular nonpeptide substance-P antagonists, which in general are metabolically more stable, and hence, may be more appropriate as pharmaceutically active substances.

[0003] Several nonpeptide tachykinin antagonists are disclosed in the art. For instance, EP-0,532,456-A, published on March 17, 1993 by Ciba-Geigy Corp., discloses 1-acylpiperidine compounds, in particular 2-arylalkyl-1-arylcarbonyl-4-piperidinamine derivatives, and their use as substance-P antagonists. EP-0,655,442-A, published on May 31, 1995 by Fujisawa Pharmaceutical Co. Ltd., discloses piperazine derivatives having tachykinin antagonistic activity.

[0004] The present compounds differ therefrom in that they invariably contain a 4-substituted(piperazine or homopiperazine)-moiety in the 4-position of a piperidine- or homopiperidine group or in the 3-position of a pyrrolidine group, and by their favourable pharmacological properties.

[0005] The present invention concerns compounds of formula



the *N*-oxide forms, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

*n* is 0, 1 or 2;

*m* is 1 or 2, provided that if *m* is 2, then *n* is 1;

*p* is 1 or 2;

*Q* is =O or =NR<sup>3</sup>;



*x* is a covalent bond or a bivalent radical of formula -O-, -S-, -NR<sup>3</sup>-;



*R*<sup>1</sup> is Ar<sup>1</sup>, Ar<sup>1</sup>C<sub>1-6</sub>alkyl or di(Ar<sup>1</sup>)C<sub>1-6</sub>alkyl, wherein each C<sub>1-6</sub>alkyl group is optionally substituted with hydroxy, C<sub>1-4</sub>alkyloxy, oxo or a ketalized oxo substituent of formula -O-CH<sub>2</sub>-CH<sub>2</sub>-O- or -O-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O-;

*R*<sup>2</sup> is Ar<sup>2</sup>, Ar<sup>2</sup>C<sub>1-6</sub>alkyl, Het<sup>1</sup> or Het<sup>1</sup>C<sub>1-6</sub>alkyl;


*R*<sup>3</sup> is hydrogen or C<sub>1-6</sub>alkyl;

*L* is hydrogen; Ar<sup>3</sup>; C<sub>1-6</sub>alkyl; C<sub>1-6</sub>alkyl substituted with 1 or 2 substituents selected from hydroxy, C<sub>1-6</sub>alkyloxy, Ar<sup>3</sup>, Ar<sup>3</sup>C<sub>1-6</sub>alkyloxy and Het<sup>2</sup>; C<sub>3-6</sub>alkenyl; Ar<sup>3</sup>C<sub>3-6</sub>alkenyl; di(Ar<sup>3</sup>)C<sub>3-6</sub>alkenyl or a radical of formula





or



wherein

- each q independently is 2, 3 or 4;  
 each r is 0, 1, 2, 3 or 4;  
 each Y<sup>1</sup> independently is a covalent bond, -O- or NR<sup>3</sup>;  
 Y<sup>2</sup> is a covalent bond, C<sub>1-4</sub>alkanediyl or -C<sub>1-4</sub>alkylNR<sup>3</sup>;  
 each -A=B- independently is a bivalent radical of formula -CH=CH-, -N=CH- or -CH=N-;  
 each R<sup>4</sup> independently is hydrogen, C<sub>1-6</sub>alkyl, Ar<sup>2</sup> or Ar<sup>2</sup>C<sub>1-6</sub>alkyl;  
 R<sup>5</sup> is hydrogen, C<sub>1-6</sub>alkyl or Ar<sup>3</sup>;  
 R<sup>6</sup> is C<sub>1-6</sub>alkyl, Ar<sup>3</sup>, Ar<sup>3</sup>C<sub>1-6</sub>alkyl, di(Ar<sup>3</sup>)C<sub>1-6</sub>alkyl, Ar<sup>3</sup>C<sub>3-7</sub>cycloalkyl, or indolyl;  
 R<sup>7</sup> is Ar<sup>3</sup>, Ar<sup>3</sup>C<sub>1-6</sub>alkyl; di(Ar<sup>3</sup>)C<sub>1-6</sub>alkyl; C<sub>1-6</sub>alkyl; C<sub>3-7</sub>cycloalkyl; C<sub>3-7</sub>cycloalkyl substituted with Ar<sup>3</sup>; oxazolyl; oxazolyl substituted with halo or C<sub>1-6</sub>alkyl; thiazolyl; thiazolyl substituted with halo or C<sub>1-6</sub>alkyl; imidazolyl; imidazolyl substituted with Ar<sup>3</sup>, C<sub>1-6</sub>alkyl, Ar<sup>3</sup>C<sub>1-6</sub>alkyl or halo; indolyl; indolyl substituted with C<sub>1-4</sub>alkyl; 2,3,4-trihydroquinolyl; pyrrolidinyl or furanyl;  
 each R<sup>8</sup> independently is hydrogen, C<sub>1-6</sub>alkyl, C<sub>3-7</sub>cycloalkyl or a radical of formula



or



wherein

- Alk is C<sub>1-6</sub>alkanediyl;  
 Z is a bivalent radical of formula -O-, -S- or -NR<sup>3</sup>;  
 R<sup>11</sup> is phenyl; phenyl substituted with 1 or 2 substituents selected from halo, C<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxy; furanyl; furanyl substituted with 1 or 2 substituents selected from halo or hydroxy; C<sub>1-6</sub>alkyl; thienyl; thienyl substituted with 1 or 2 substituents selected from halo or C<sub>1-6</sub>alkyl; oxazolyl; oxazolyl substituted with 1 or 2 C<sub>1-6</sub>alkyl substituents; thiazolyl; thiazolyl substituted with 1 or 2 C<sub>1-6</sub>alkyl substituents; pyridinyl or pyridinyl substituted with 1 or 2 C<sub>1-6</sub>alkyl substituents;

R<sup>12</sup> is C<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyl substituted with hydroxy, carboxyl or C<sub>1-6</sub>alkyloxycarbonyl;

Ar<sup>1</sup> is phenyl; phenyl substituted with 1, 2 or 3 substituents each independently selected from halo, C<sub>1-4</sub>alkyl, haloC<sub>1-4</sub>alkyl, cyano, aminocarbonyl, C<sub>1-4</sub>alkyloxy or haloC<sub>1-4</sub>alkyloxy;

Ar<sup>2</sup> is naphthalenyl; phenyl; phenyl substituted with 1, 2 or 3 substituents each independently selected from hydroxy, halo, cyano, nitro, amino, mono- or di(C<sub>1-4</sub>alkyl)amino, C<sub>1-4</sub>alkyl, haloC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyloxy, haloC<sub>1-4</sub>alkyloxy, carboxyl, C<sub>1-4</sub>alkyloxycarbonyl, aminocarbonyl and mono- or di(C<sub>1-4</sub>alkyl)aminocarbonyl;

Ar<sup>3</sup> is phenyl or phenyl substituted with 1, 2 or 3 substituents selected from halo, hydroxy, amino, nitro, aminocarbonyl, C<sub>1-6</sub>alkyl, haloC<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxy;

Het<sup>1</sup> is a monocyclic heterocycle selected from pyrrolyl, pyrazolyl, imidazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl; or a bicyclic heterocycle selected from quinolinyl, quinoxalinyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl and benzothienyl; each monocyclic and bicyclic heterocycle may optionally be substituted on a carbon atom by 1 or 2 substituents selected from halo, C<sub>1-4</sub>alkyl or mono-, di- or tri(halo)methyl; and

Het<sup>2</sup> is a heterocycle selected from 1,4-dihydro-5-oxo-tetrazol-1-yl, imidazo[1,2-a]-pyridinyl, oxazolyl or imidazolyl; each of said heterocycles may be substituted with 1 or where possible 2 substituents selected from C<sub>1-4</sub>alkyl and Ar<sup>3</sup>.

[0006] The heterocycles in the definition of Het<sup>1</sup> are preferably connected to the rest of the molecule, i.e. X, -C(=Q)- or C<sub>1-6</sub>alkyl, by a carbon atom.

[0007] As used in the foregoing definitions and hereinafter, halo is generic to fluoro, chloro, bromo and iodo; C<sub>2-4</sub>alkyl defines straight and branched chain saturated hydrocarbon radicals having from 2 to 4 carbon atoms such as, for example, ethyl, propyl, butyl, 1-methylethyl, 2-methylpropyl and the like; C<sub>1-4</sub>alkyl is meant to include C<sub>2-4</sub>alkyl and methyl; C<sub>1-5</sub>alkyl is meant to include C<sub>1-4</sub>alkyl and the higher homologues thereof having 5 carbon atoms such as, for example, pentyl, 2-methylbutyl and the like; C<sub>1-6</sub>alkyl is meant to include C<sub>1-5</sub>alkyl and the higher homologues thereof having 6 carbon atoms such as, for example, hexyl, 2-methylpentyl and the like; C<sub>1-4</sub>alkanediyl defines bivalent straight and branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as, for example, methylene, 1,2-ethanediyl, 1,3-propanediyl, 1,4-butanediyl, and the like; C<sub>1-6</sub>alkanediyl is meant to include C<sub>1-4</sub>alkanediyl and the higher homologues thereof having from 5 to 6 carbon atoms such as, for example, 1,5-pentanediyl, 1,6-hexanediyl and the like; C<sub>3-6</sub>alkenyl defines straight and branched chain hydrocarbon radicals containing one double bond and having from 3 to 6 carbon atoms such as, for example, 2-propenyl, 3-butenyl, 2-butenyl, 2-pentenyl, 3-pentenyl, 3-methyl-2-butenyl, 3-hexenyl and the like; and the carbon of said C<sub>3-6</sub>alkenyl connected to the nitrogen atom of the piperazine or homopiperazine preferably is saturated.

[0008] As used in the foregoing definitions and hereinafter, haloC<sub>1-4</sub>alkyl is defined as mono- or polyhalosubstituted C<sub>1-4</sub>alkyl, in particular C<sub>1-4</sub>alkyl substituted with 1 to 6 halogen atoms, more in particular difluoro- or trifluoromethyl.

[0009] The pharmaceutically acceptable addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid addition salt forms which the compounds of formula (I) are able to form. Said salts can conveniently be obtained by treating the base form of the compounds of formula (I) with appropriate acids such as, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid; sulfuric; nitric; phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pantoic and the like acids.

[0010] The pharmaceutically acceptable addition salts as mentioned hereinabove are also meant to comprise the therapeutically active non-toxic base, in particular, a metal or amine addition salt forms which the compounds of formula (I) are able to form. Said salts can conveniently be obtained by treating the compounds of formula (I) containing acidic hydrogen atoms with appropriate organic and inorganic bases such as, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, *N*-methyl-D-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

[0011] Conversely said salt forms can be converted by treatment with an appropriate base or acid into the free acid or base form.

[0012] The term addition salt as used hereinabove also comprises the solvates which the compounds of formula (I) as well as the salts thereof, are able to form. Such solvates are for example hydrates, alcoholates and the like.

[0013] For isolation and purification purposes, it is also possible to use pharmaceutically unacceptable salts. Only the pharmaceutically acceptable, non-toxic salts are used therapeutically and those salts are therefore preferred.

[0014] The term "stereochemically isomeric forms" as used hereinbefore defines all the possible isomeric as well as conformational forms which the compounds of formula (I) may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture, more in particular the racemic mixture, of all possible ster-

eochemically and conformationally isomeric forms, said mixtures containing all diastereomers, enantiomers and/or conformers of the basic molecular structure. More in particular, stereogenic centers may have the R- or S-configuration; substituents on bivalent cyclic saturated radicals may have either the *cis*- or *trans*-configuration;  $>C=NR^3$  and  $C_{3-6}$ alkenyl radicals may have the E- or Z-configuration. The compounds of formula (I) have at least two stereogenic centers; thus for compounds of which the actual stereochemical configuration is known, the relative stereodescriptors R\* and S\* may be used in accordance with the Chemical Abstracts rules (Chemical Substance Name Selection Manual (CA), 1982 Edition, Vol. III, Chapter 20). In those cases where the compounds of formula (I) were separated into its racemic *cis* and racemic *trans* isomers, or in those cases where the racemic *cis* or racemic *trans* isomers were separated into its pure enantiomeric forms, the stereochemically isomeric form which was first isolated was designated as "A" and the second as "B". All stereochemically isomeric forms of the compounds of formula (I) both in pure form or mixtures thereof are intended to be embraced within the scope of the present invention.

[0015] Some of the compounds of formula (I) may also exist in their tautomeric form. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention. For instance, compounds of formula (I) wherein L is a radical of formula (a-1) wherein  $R^5$  is hydrogen, or a radical of (a-2) or (a-3) wherein  $Y^1$  is -NH-, or a radical of formula (a-5) wherein  $R^3$  is hydrogen may exist in their corresponding tautomeric form. Also compounds of formula (I) wherein X is -NH- and =Q is =O may exist in their corresponding tautomeric form.

[0016] The N-oxide forms of the compounds of formula (I) are meant to comprise those compounds of formula (I) wherein one or several nitrogen atoms are oxidized to the so-called N-oxide, particularly those N-oxides wherein one or more of the piperazine-nitrogens are N-oxidized.

[0017] Whenever used hereinafter, the term "compounds of formula (I)" is meant to also include their N-oxide forms, their pharmaceutically acceptable addition salts, and their stereochemically isomeric forms.

[0018] A special group of compounds are those compounds of formula (I) wherein L is hydrogen;  $C_{1-6}$ alkyl;  $C_{1-6}$ alkyl substituted with hydroxy;  $C_{3-6}$ alkenyl;  $Ar^3$ ;  $Ar^3C_{1-6}$ alkyl;  $di(Ar^3)C_{1-6}$ alkyl;  $Ar^3C_{3-6}$ alkenyl;  $di(Ar^3)C_{1-6}$ alkenyl; or a radical of formula (a-1), (a-2), (a-4) or (a-5) wherein

$R^7$  is  $Ar^3$ ;  $Ar^3C_{1-6}$ alkyl;  $di(Ar^3)C_{1-6}$ alkyl;  $C_{1-6}$ alkyl;  $C_{3-7}$ cycloalkyl;  $C_{3-7}$ cycloalkyl substituted with  $Ar^3$ ; oxazolyl; oxazolyl substituted with halo or  $C_{1-6}$ alkyl; thiazolyl; thiazolyl substituted with halo or  $C_{1-6}$ alkyl; imidazolyl; imidazolyl substituted with  $Ar^3$ ,  $C_{1-6}$ alkyl,  $Ar^3C_{1-6}$ alkyl or halo; pyrrolidinyl or furanyl;

$Ar^3$  is phenyl or phenyl substituted with 1, 2 or 3 substituents selected from halo, hydroxy, amino, aminocarbonyl,  $C_{1-6}$ alkyl, halo $C_{1-6}$ alkyl or  $C_{1-6}$ alkyloxy;

Het<sup>1</sup> is a monocyclic heterocycle selected from pyrrolyl, pyrazolyl, imidazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl; or a bicyclic heterocycle selected from quinolinyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl and benzothienyl; each monocyclic and bicyclic heterocycle may optionally be substituted on a carbon atom by 1 or 2 substituents selected from halo,  $C_{1-4}$ alkyl or mono-, di- or tri(halo)methyl.

[0019] A first group of interesting compounds consists of those compounds of formula (I) wherein one or more of the following restrictions apply :

a)  $R^1$  is  $Ar^1C_{1-6}$ alkyl; or

b)  $R^2$  is  $Ar^2$ ,  $Ar^2C_{1-6}$ alkyl or Het<sup>1</sup>; in particular, phenyl substituted with 1, 2 or 3 substituents each independently selected from halo, cyano, nitro, amino,  $C_{1-4}$ alkyl, halo $C_{1-4}$ alkyl,  $C_{1-4}$ alkyloxy and  $C_{1-4}$ alkyloxycarbonyl, more in particular, phenyl substituted with 2 substituents selected from methyl and trifluoromethyl; or

c) n is 0 or 1, in particular n is 1; or

d) m is 1; or

e) p is 1 or 2, in particular p is 1; or

f) =Q is =O; or

g) X is a covalent bond, -O- or -NR<sup>3</sup>-, in particular a covalent bond.

[0020] A second group of interesting compounds consists of those compounds of formula (I) wherein L is hydrogen,  $Ar^3$ ;  $Ar^3C_{1-6}$ alkyl;  $di(Ar^3)C_{1-6}$ alkyl;  $Ar^3C_{3-6}$ alkenyl;  $C_{1-6}$ alkyl substituted with hydroxy; or

a radical of formula (a-2) wherein

R<sup>4</sup> is hydrogen or Ar<sup>2</sup>;

r is 0 or 1;

Y<sup>1</sup> is a covalent bond, -O- or -NR<sup>3</sup>-; and

R<sup>7</sup> is Ar<sup>3</sup>, C<sub>3-7</sub>cycloalkyl substituted with Ar<sup>3</sup>, di(Ar<sup>3</sup>)methyl, pyrrolidinyl or furanyl; or

a radical of formula (a-4) wherein

Y<sup>2</sup> is a covalent bond or methylene;

-A=B- is -CH=CH- or -N=CH-; and

R<sup>8</sup> is hydrogen, a radical of formula (b-1) wherein R<sup>11</sup> is methyl substituted oxazolyl, or a radical of formula (b-2) wherein Z is -O- and R<sup>12</sup> is C<sub>1-6</sub>alkyl; or

a radical of formula (a-5) wherein

R<sup>4</sup> is hydrogen;

q is 2; and

R<sup>3</sup> is hydrogen.

[0021] A third group of interesting compounds consists of those compounds of formula (I) wherein

q is 2 or 4;

-A=B- is -CH=CH- or -N=CH-;

R<sup>4</sup> is hydrogen or Ar<sup>2</sup>;

R<sup>5</sup> is hydrogen;

R<sup>6</sup> is C<sub>1-6</sub>alkyl or Ar<sup>3</sup>;

R<sup>7</sup> is Ar<sup>3</sup>, di(Ar<sup>3</sup>)C<sub>1-6</sub>alkyl; C<sub>1-6</sub>alkyl; C<sub>3-7</sub>cycloalkyl substituted with Ar<sup>3</sup>; thiazolyl; imidazolyl substituted with C<sub>1-6</sub>alkyl or Ar<sup>3</sup>C<sub>1-6</sub>alkyl; indolyl; indolyl substituted with C<sub>1-4</sub>alkyl; 2,3,4-trihydroquinolyl; pyrrolidinyl or furanyl;

Z is -O-;

R<sup>11</sup> is phenyl substituted with halo; oxazolyl substituted with C<sub>1-6</sub>alkyl; or

R<sup>12</sup> is C<sub>1-6</sub>alkyl.

[0022] Of special interest are those compounds of formula (I) wherein R<sup>1</sup> is Ar<sup>1</sup>C<sub>1-6</sub>alkyl, R<sup>2</sup> is phenyl substituted with 2 substituents selected from methyl or trifluoromethyl, X is a covalent bond and =Q is =O.

[0023] Further of special interest are those compounds of formula (I) wherein n and m are 1 and p is 1 or 2.

[0024] Particular compounds are those compounds of formula (I) wherein

R<sup>1</sup> is phenylmethyl;

R<sup>2</sup> is phenyl substituted with 2 substituents selected from methyl or trifluoromethyl;

n, m and p are 1;

X is a covalent bond; and

=Q is =O.

[0025] Also particular compounds are those compounds of formula (I) wherein L is a radical of formula (a-2) wherein

R<sup>4</sup> is hydrogen or phenyl;

r is 0 or 1;

Y<sup>1</sup> is a covalent bond, -O- or -NH-;

R<sup>7</sup> is pyrrolidinyl; furanyl; 1-phenylcyclohexanyl; diphenylmethyl; or phenyl substituted with 1, 2 or 3 substituents each independently selected from methyl, methoxy or chloro.

[0026] Preferred compounds are those particular compounds that have a *trans* configuration.

[0027] Other preferred compounds are those particular compounds that have a *cis* configuration.

[0028] Still other preferred compounds are those compounds of formula (I) wherein

R<sup>1</sup> is phenylmethyl;

$R^2$  is phenyl substituted with 2 substituents selected from methyl or trifluoromethyl;  
 $n, m$  and  $p$  are 1;  
 $x$  is a covalent bond;  
 $=Q$  is =O;  
 $L$  is a radical of formula (a-2) wherein

$R^4$  is hydrogen;  
 $r$  is 1;  
 $Y^1$  is -NH-; and  
 $R^7$  is phenyl substituted with 2 methyl substituents.


[0029] Most preferred are those compounds selected from

4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidiny]-*N*-(2,6-dimethylphenyl)-1-piperazine acetamide;  
 4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidiny]-*N*-(1-phenylcyclohexyl)-1-piperazine acetamide;  
 1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-[4-[ $\alpha$ -(1-pyrrolidinylcarbonyl)benzyl]-1-piperazinyl]piperidine;  
 1-[3,5-bis(trifluoromethyl)benzoyl]-4-[4-[1-[(2-methyl-5-oxazolyl)methyl]-1*H*-benzimidazol-2-yl]-1-piperazinyl]-2-(phenylmethyl)piperidine;  
 4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(4-trifluoromethylphenyl)methyl]-4-piperidiny]-*N*-(2,6-dimethylphenyl)-1-piperazine acetamide;  
 4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-[(3,4-dichlorophenyl)methyl]-4-piperidiny]-*N*-(2,6-dimethylphenyl)-1-piperazine acetamide; the *N*-oxides, the stereoisomeric forms and the pharmaceutically acceptable addition salts thereof.

[0030] Particularly interesting stereoisomeric forms are

(+)-(*B*)-*trans*-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidiny]-*N*-(2,6-dimethylphenyl)-1-piperazine acetamide; and  
 (-)-(*B*)-*cis*-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidiny]-*N*-(2,6-dimethylphenyl)-1-piperazine acetamide, and the pharmaceutically acceptable addition salts thereof, especially the (*L*) malic acid form.


[0031] The compounds of formula (I) can be prepared by reductively *N*-alkylating an intermediate of formula (III) with an intermediate of formula (II). Said reductive *N*-alkylation may be performed in a reaction-inert solvent such as, for example, dichloromethane, ethanol, toluene or a mixture thereof, and in the presence of an appropriate reducing agent such as, for example, a borohydride, e.g. sodium borohydride, sodium cyanoborohydride or triacetoxy borohydride. In case a borohydride is used as a reducing agent, it may be convenient to use a complex-forming agent such as, for example, titanium(IV)isopropylate as described in J. Org. Chem, 1990, 55, 2552-2554. Using said complex-forming agent may also result in an improved *cis/trans* ratio in favour of the *trans* isomer. It may also be convenient to use hydrogen as a reducing agent in combination with a suitable catalyst such as, for example, palladium-on-charcoal or platinum-on-charcoal. In case hydrogen is used as reducing agent, it may be advantageous to add a dehydrating agent to the reaction mixture such as, for example, aluminium *tert*-butoxide. In order to prevent the undesired further hydrogenation of certain functional groups in the reactants and the reaction products, it may also be advantageous to add an appropriate catalyst-poison to the reaction mixture, e.g., thiophene or quinoline-sulphur. Stirring and optionally elevated temperatures and/or pressure may enhance the rate of the reaction.




[0032] In this and the following preparations, the reaction products may be isolated from the reaction medium and,

if necessary, further purified according to methodologies generally known in the art such as, for example, extraction, crystallization, trituration and chromatography.


[0033] The compounds of formula (I) can also be prepared by reacting an intermediate of formula (IV) wherein W<sup>1</sup> is an appropriate leaving group such as, for example, a halogen, e.g. chloro or bromo, or a sulfonyloxy leaving group, e.g. methanesulfonyloxy or benzenesulfonyloxy, with an intermediate of formula (V). The reaction can be performed in a reaction-inert solvent such as, for example, a chlorinated hydrocarbon, e.g. dichloromethane, an alcohol, e.g. ethanol, or a ketone, e.g. methyl isobutylketone, and in the presence of a suitable base such as, for example, sodium carbonate, sodium hydrogen carbonate or triethylamine. Stirring may enhance the rate of the reaction. The reaction may conveniently be carried at a temperature ranging between room temperature and reflux temperature.



[0034] The compounds of formula (I) may also be converted into each other following art-known transformations. In particular, the compounds of formula (I) wherein L is other than hydrogen, said L being represented by L' and said compounds being represented by formula (I-a), can also be prepared by reacting a compound of formula (I) wherein L is hydrogen, said compounds being represented by formula (I-b), with an intermediate of formula (VI) wherein W<sup>2</sup> is an appropriate leaving group such as, for example, a halogen, e.g. chloro or bromo, or a sulfonyloxy leaving group, e.g. methanesulfonyloxy or benzenesulfonyloxy, at reaction conditions which are similar to those for the reaction between intermediates of formula (IV) and (V).




[0035] Compounds of formula (I-b) may be prepared by reductively N-alkylating a piperazine derivative of formula (VII) wherein P<sup>1</sup> is a protective group such as, for example, benzyl, with an intermediate of formula (II). Said reaction may be performed in a similar way as described hereinabove for the reductive N-alkylation using intermediates (II) and (III). The thus formed compound of formula (I-c) may then be deprotected using art-known deprotection techniques. Depending on the nature of the protective group P<sup>1</sup>, compounds of formula (I-c) may be part of the scope of the compounds of formula (I).





[0036] Alternatively, compounds of formula (I-b) may be prepared by first reductively *N*-alkylating a piperazine derivative of formula (VII) wherein P<sup>1</sup> is a protective group such as, for example, halo, with an intermediate of formula (VIII) using the same procedure as described hereinabove for the reductive *N*-alkylation using intermediates (II) and (III). The thus formed intermediate of formula (XI) may then be reacted with an intermediate of formula (IV) in a reaction-inert solvent and optionally in the presence of a suitable base such as, for example, triethylamine, to form a compound of formula (I-c), which may then be deprotected using art-known deprotection techniques.




[0037] The compounds of formula (I-b) are deemed to be of particular use in the synthesis of other compounds of formula (I).

[0038] The compounds of formula (I) may also be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboperoxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chlorobenzenecarboperoxoic acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. *tert*-butyl hydroperoxide. Suitable solvents are, for example, water, lower alkanols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.


[0039] The starting materials and some of the intermediates are known compounds and are commercially available or may be prepared according to conventional reaction procedures generally known in the art. For example, intermediates of formula (III), (IV) and (VI) may be prepared according to art-known procedures.

[0040] Intermediates of formula (II) may be prepared by condensing an intermediate of formula (IV) with an intermediate of formula (VIII) analogous to the procedure described in EP-0,532,456-A.




[0041] The preparation of intermediates of formula (VIII) is also described in EP-0,532,456-A. However, intermediates of formula (VIII) wherein R<sup>1</sup> is optionally substituted Ar<sup>1</sup>C<sub>1-6</sub>alkyl or di(Ar<sup>1</sup>)C<sub>1-6</sub>alkyl, said R<sup>1</sup> being represented by -CH(R<sup>1a</sup>)<sub>2</sub> and said intermediates being represented by formula (VIII-a), may also be prepared as depicted in scheme 1.

## Scheme 1




[0042] In scheme 1, the intermediates of formula (IX-b) may be prepared by reacting an intermediate of formula (IX-a) with an aldehyde or a ketone of formula (X). The  $\text{C}_{1-6}$ alkylcarbamate moiety in the intermediates of formula (IX-b) may be converted into a fused oxazolone which in turn may be reduced to an intermediate of formula (IX-d). Said intermediate (IX-d) may in turn be deprotected, thus forming an intermediate of formula (VIII-a). Subsequently, intermediates of formula (VIII-a) may be reacted with an intermediate of formula (IV) to prepare intermediates of formula (II) wherein  $\text{R}^1$  is defined as  $-\text{CH}(\text{R}^{1a})_2$ , said intermediates being represented by formula (II-a).

[0043] Said intermediates of formula (II-a) may also be prepared by first reacting intermediate (IX-d) with intermediate (IV) in the presence of a suitable base to form an intermediate of formula (XII), which may subsequently be deprotected. These reactions and those performed in scheme 1 may all be conducted following conventional methods that are generally known in the art.




Intermediates of formula (V) may suitably be prepared by reacting an intermediate of formula (VIII-1), being a protected intermediate of formula (VIII) with a protecting group  $\text{P}^2$  such as, for example, a  $\text{C}_{1-6}$ alkyloxycarbonyl group, with an intermediate of formula (III) according to the previously described reductive *N*-alkylation procedure, and subsequently deprotecting the thus formed intermediate.



[0044] In particular, intermediates of formula (V) wherein  $\text{R}^1$  is  $-\text{CH}(\text{R}^{1a})_2$ , said intermediates being represented by formula (V-a), may be prepared as is depicted in scheme 2.

### Scheme 2



[0045] The ketalized intermediate of formula (IX-c) may be transformed to the corresponding ketone of formula (IX-e) which subsequently may be reductively aminated with a piperazine- or homopiperazine derivative of formula (III).

The thus obtained intermediate may then be reduced with a suitable reducing agent to an intermediate of formula (V-a). [0046] Pure stereochemically isomeric forms of the compounds of formula (I) may be obtained by the application of art-known procedures. Diastereomers may be separated by physical methods such as selective crystallization and chromatographic techniques, e.g., countercurrent distribution, liquid chromatography and the like.

[0047] The compounds of formula (I) as prepared in the hereinabove described processes are generally racemic mixtures of enantiomers which can be separated from one another following art-known resolution procedures. The racemic compounds of formula (I) which are sufficiently basic or acidic may be converted into the corresponding diastereomeric salt forms by reaction with a suitable chiral acid, respectively chiral base. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali or acid. An alternative manner of separating the enantiomeric forms of the compounds of formula (I) involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

[0048] The compounds of formula (I) have valuable pharmacological properties in that they interact with tachykinin receptors and they antagonize tachykinin-induced effects, especially substance P-induced effects, both *in vivo* and *in vitro* and are thus of use in the treatment of tachykinin-mediated diseases, and in particular in substance P-mediated diseases.

[0049] Tachykinins, also referred to as neurokinins, are a family of peptides among which substance P (SP), neurokinin A (NKA), neurokinin B (NKB) and neuropeptide K (NPK) may be identified. They are naturally occurring in mammals, including human beings, and are distributed throughout the central and peripheral nervous system, where they act as neurotransmitters or neuromodulators. Their actions are mediated through several subtypes of receptors, such

as, for example, NK<sub>1</sub>, NK<sub>2</sub> and NK<sub>3</sub> receptors. Substance P displays highest affinity for NK<sub>1</sub> receptors, whereas NKA preferentially binds to NK<sub>2</sub> receptors and NKB preferentially binds to NK<sub>3</sub> receptors. However, the selectivity of these tachykinins is relatively poor and under physiological conditions the action of any of these tachykinins might be mediated by activation of more than one receptor type.

[0050] Substance P and other neurokinins are involved in a variety of biological actions such as pain transmission (nociception), neurogenic inflammation, smooth muscle contraction, plasma protein extravasation, vasodilation, secretion, mast cell degranulation, and also in activation of the immune system. A number of diseases are deemed to be engendered by activation of neurokinin receptors, in particular the NK<sub>1</sub> receptor, by excessive release of substance P and other neurokinins in particular cells such as cells in the neuronal plexi of the gastrointestinal tract, unmyelinated primary sensory afferent neurons, sympathetic and parasympathetic neurons and nonneuronal cell types (DN&P 8(1), February 1995, p. 5-23, "Neurokinin Receptors" by Longmore J. et al.; Pharmacological Reviews 46(4), 1994, p. 551-599, "Receptors and Antagonists for Substance P and Related Peptides" by Regoli et al.).

[0051] The compounds of the present invention are potent inhibitors of neurokinin-mediated effects, in particular those mediated via the NK<sub>1</sub> receptor, and may therefore be described as tachykinin antagonists, especially as substance P antagonists, as indicated *in vitro* by the antagonism of substance P-induced relaxation of pig coronary arteries which is described hereinafter. The binding affinity of the present compounds for the human, guinea-pig and gerbil neurokinin receptors may be determined *in vitro* in a receptor binding test using <sup>3</sup>H-substance-P as radioligand. The subject compounds also show substance-P antagonistic activity *in vivo* as may be evidenced by, for instance, the antagonism of substance P-induced plasma extravasation in guinea-pigs, or the antagonism of drug-induced emesis in ferrets (Watson et al., Br. J. Pharmacol. 115, 84-94, 1995).

[0052] In view of their capability to antagonize the actions of tachykinins by blocking the tachykinin receptors, and in particular antagonizing the actions of substance P by blocking the NK<sub>1</sub> receptor, the subject compounds are useful in the prophylactic and therapeutic treatment of tachykinin-mediated diseases such as, for example,

- pain, in particular traumatic pain such as postoperative pain; traumatic avulsion pain such as brachial plexus; chronic pain such as arthritic pain such as occurring in osteo-, rheumatoid or psoriatic arthritis; neuropathic pain such as post-herpetic neuralgia, trigeminal neuralgia, segmental or intercostal neuralgia, fibromyalgia, causalgia, peripheral neuropathy, diabetic neuropathy, chemotherapy-induced neuropathy, AIDS-related neuropathy, occipital neuralgia, geniculate neuralgia, glossopharyngeal neuralgia, reflex sympathetic dystrophy, phantom limb pain; various forms of headache such as migraine, acute or chronic tension headache, temperomandibular pain, maxillary sinus pain, cluster headache; odontalgia; cancer pain; pain of visceral origin; gastrointestinal pain; nerve entrapment pain; sport's injury pain; dysmenorrhoea; menstrual pain; meningitis; arachnoiditis; musculoskeletal pain; low back pain e.g. spinal stenosis; prolapsed disc; sciatica; angina; ankylosing spondylitis; gout; burns; scar pain; itch; and thalamic pain such as post stroke thalamic pain;
- respiratory and inflammatory diseases, in particular inflammation in asthma, influenza, chronic bronchitis and rheumatoid arthritis; inflammatory diseases of the gastrointestinal tract such as Crohn's disease, ulcerative colitis, inflammatory bowel disease and non-steroidal anti-inflammatory drug induced damage; inflammatory diseases of the skin such as herpes and eczema; inflammatory diseases of the bladder such as cystitis and urge incontinence; and eye and dental inflammation;
- emesis, i.e. nausea, retching and vomiting, including acute emesis, delayed emesis and anticipatory emesis, no matter how emesis is induced, for example, emesis may be induced by drugs such as cancer chemotherapeutic agents such as alkylating agents, e.g. cyclophosphamide, carmustine, lomustine and chlorambucil; cytotoxic antibiotics, e.g. dactinomycin, doxorubicin, mitomycin-C and bleomycin; antimetabolites, e.g. cytarabine, methotrexate and 5-fluorouracil; vinca alkaloids, e.g. etoposide, vinblastine and vincristine; and others such as cisplatin, dacarbazine, procarbazine and hydroxyurea; and combinations thereof; radiation sickness; radiation therapy, e.g. irradiation of the thorax or abdomen, such as in the treatment of cancer; poisons; toxins such as toxins caused by metabolic disorders or by infection, e.g. gastritis, or released during bacterial or viral gastrointestinal infection; pregnancy; vestibular disorders, such as motion sickness, vertigo, dizziness and Ménière's disease; post-operative sickness; gastrointestinal obstruction; reduced gastrointestinal motility; visceral pain, e.g. myocardial infarction or peritonitis; migraine; increased intracranial pressure; decreased intracranial pressure (e.g. altitude sickness); opioid analgesics, such as morphine; and gastro-oesophageal reflux disease, acid indigestion, over-indulgence of food or drink, acid stomach, sour stomach, waterbrash/regurgitation, heartburn, such as episodic heartburn, nocturnal heartburn, and meal-induced heartburn and dyspepsia;
- central nervous system disorders, in particular psychoses such as schizophrenia, mania, dementia or other cognitive disorders e.g. Alzheimer's disease; anxiety; AIDS-related dementia; diabetic neuropathy; multiple sclerosis; depression; Parkinson's disease; and dependence on drugs or substances of abuse;
- allergic disorders, in particular allergic disorders of the skin such as urticaria, and allergic disorders of the airways such as rhinitis;

- gastrointestinal disorders, such as irritable bowel syndrome;
- skin disorders, such as psoriasis, pruritis and sunburn;
- vasospastic diseases, such as angina, vascular headache and Reynaud's disease;
- cerebral ischaemia, such as cerebral vasospasm following subarachnoid haemorrhage;
- 5 - stroke, epilepsie, head trauma, spinal cord trauma and ischemic neuronal damage;
- fibrosing and collagen diseases, such as scleroderma and eosinophilic fascioliasis;
- disorders related to immune enhancement or suppression, such as systemic lupus erythematosus;
- rheumatic diseases, such as fibrositis;
- neoplastic disorders;
- 10 - cell proliferation; and
- cough.

[0053] The compounds of the present invention have a favourable metabolic stability and exhibit good oral availability. They also have an advantageous onset and duration of action. The compounds of formula (I) also have the ability to  
15 penetrate the central nervous system as may be demonstrated *in vivo* by their inhibitory effect on the change in behaviour induced by intracerebroventricular-applied substance P in the gerbil.

[0054] In view of the utility of the compounds of formula (I), there is provided a method of treating warm-blooded animals, including humans, suffering from tachykinin-mediated diseases as mentioned hereinabove, in particular, pain, emesis or asthma. Said method comprises the systemic administration of an effective tachykinin antagonizing amount  
20 of a compound of formula (I), a *N*-oxide form, a pharmaceutically acceptable addition salt or a possible stereoisomeric form thereof, to warm-blooded animals, including humans. Hence, the use of a compound of formula (I) as a medicine is provided, and in particular a medicine to treat pain, emesis or asthma.

[0055] For ease of administration, the subject compounds may be formulated into various pharmaceutical forms for administration purposes. To prepare the pharmaceutical compositions of this invention, a therapeutically effective  
25 amount of the particular compound, optionally in addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for  
30 example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though  
35 other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable solutions containing compounds of formula (I) may be formulated in an oil for prolonged action. Appropriate oils for this purpose are, for example, peanut oil, sesame oil, cottonseed oil, corn oil, soy bean oil, synthetic glycerol esters of long chain fatty acids and mixtures of these and other oils. Injectable suspensions may also be prepared in which case  
40 appropriate liquid carriers, suspending agents and the like may be employed. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wettable agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not cause any significant deleterious effects on the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as  
45 a transdermal patch, as a spot-on or as an ointment. Acid or base addition salts of compounds of formula (I) due to their increased water solubility over the corresponding base or acid form, are obviously more suitable in the preparation of aqueous compositions.

[0056] In order to enhance the solubility and/or the stability of the compounds of formula (I) in pharmaceutical compositions, it can be advantageous to employ  $\alpha$ -,  $\beta$ - or  $\gamma$ -cyclodextrins or their derivatives, in particular hydroxyalkyl  
50 substituted cyclodextrins, e.g. 2-hydroxypropyl- $\beta$ -cyclodextrin. Also co-solvents such as alcohols may improve the solubility and/or the stability of the compounds of formula (I) in pharmaceutical compositions.

[0057] It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims  
55 herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect, in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

[0058] Those of skill in the treatment of tachykinin mediated diseases could determine the effective therapeutic daily amount from the test results presented hereinafter. An effective therapeutic daily amount would be from about 0.001 mg/kg to about 40 mg/kg body weight, more preferably from about 0.01 mg/kg to about 5 mg/kg body weight. It may be appropriate to administer the therapeutically effective dose once daily or as two, three, four or more sub-doses at appropriate intervals throughout the day. Said sub-doses may be formulated as unit dosage forms, for example, containing 0.05 mg to 500 mg, and in particular, 0.5 mg to 50 mg of active ingredient per unit dosage form.

[0059] The exact dosage and frequency of administration depends on the particular compound of formula (I) used, the particular condition being treated, the severity of the condition being treated, the age, weight and general physical condition of the particular patient as well as other medication the patient may be taking, as is well known to those skilled in the art. Furthermore, it is evident that said effective daily amount may be lowered or increased depending on the response of the treated patient and/or depending on the evaluation of the physician prescribing the compounds of the instant invention. The effective daily amount ranges mentioned hereinabove are therefore only guidelines.

[0060] The following examples are intended to illustrate and not to limit the scope of the present invention.

## Experimental Part

[0061] Hereinafter "RT" means room temperature, "THF" means tetrahydrofuran, "DIPE" means diisopropylether, "DCM" means dichloromethane and "DMF" means *N,N*-dimethylformamide.

### A. Preparation of the intermediate compounds

#### Example A.1

[0062]

a) A mixture of ( $\pm$ )-1,1-dimethyl 7-(phenylmethyl)-1,4-dioxo-8-azaspiro[4,5]decane-8-carboxylate (13 g; prepared according to the method described in EP-A-532,456) in HCl (6N; 130 ml) was stirred and refluxed for 3 hours. The reaction mixture was cooled, alkalinized with aqueous NaOH (50 %) and extracted with DCM. The organic layer was separated, dried, filtered, and the filtrate, which contained ( $\pm$ )-2-(phenylmethyl)-4-piperidinone (intermediate 1), was used in next reaction step.

b) A mixture of the filtrate obtained in the previous reaction step, 3,5-dimethylbenzoyl chloride (7.4 g) and triethylamine (11 ml) was stirred overnight at RT. The reaction mixture was extracted with dilute NaOH solution. The organic layer was separated, dried, filtered and the solvent evaporated. The residue was crystallized from DIPE. The precipitate was filtered off and dried, yielding 7.44 g (58%) of ( $\pm$ )-1-(3,5-dimethylbenzoyl)-2-(phenylmethyl)-4-piperidinone (intermediate 2; mp. 107.8°C).

#### Example A.2

[0063] a) A mixture of ( $\pm$ )-1,1-dimethyl 7-(phenylmethyl)-1,4-dioxo-8-azaspiro[4,5]decane-8-carboxylate (33.34 g; prepared according to the method described in EP-A-532,456) in HCl (6N; 250 ml) was stirred at 70 °C for 1 hour and 30 minutes. The mixture was cooled, alkalinized with NaOH while cooling to 25°C, and extracted with DCM (100 ml). The organic layer was separated and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>. Triethylamine (20.2 g), followed by 3,5-bis(trifluoromethyl)benzoyl chloride (27.7 g) dissolved in a little DCM were added and the mixture was stirred for 2 hours. The mixture was extracted with water, and the layers were separated. The organic layer was dried, filtered and the solvent evaporated. The residue was crystallized from DIPE, the precipitate was filtered off and dried, yielding 18.34 g product. The mother layer was evaporated and the residue was crystallized from DIPE. The precipitate was filtered off and dried, yielding 6.51 g of product. The two fractions were put together and taken up in water and DCM, NaOH was added and the mixture was extracted. The organic layer was dried, filtered and the solvent evaporated, yielding 16.14 g (38%) of ( $\pm$ )-1-[3,5-bis-(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinone (intermediate 3; mp. 102.5°C).

#### Example A.3

[0064] A mixture of pyrrolidine (2.13 g) and triethylamine (6.06 g) in DCM (100 ml) was stirred at -10°C. 2-chloro-2-phenylacetylchloride (5.67 g) was added slowly and dropwise. The mixture was allowed to warm to RT and was then stirred overnight. The mixture was extracted with water and K<sub>2</sub>CO<sub>3</sub>. The separated organic layer was dried, filtered and the solvent was evaporated. The residue was crystallized from DIPE and the precipitate was filtered off and dried, yielding 3.25 g (48 %) of fraction 1. The mother layer was separated and the solvent was evaporated. The residue was

crystallized from DIPE and the precipitate was filtered off and dried, yielding 0.29 g (5 %) of fraction 2. Both fractions were combined, thus yielding 3.54 g (53 %) of ( $\pm$ )-1-(2-chloro-2-phenylacetyl)pyrrolidine (intermediate 4; mp. 88.5 °C).

#### Example A.4

[0065] Sodium hydride (2 g) was added portion wise to a solution of 3,5-dimethylphenol (6.1 g) in DMF (50 ml). The mixture was stirred for 30 minutes and added dropwise at a temperature below 30 °C to a solution of 2-chloro-2-phenylacetylchloride (9.45 g) in DMF (50 ml). The mixture was stirred overnight, decomposed with water (5 ml) and the solvent was evaporated. Water was added and the mixture was extracted with DCM. The separated organic layer was dried, filtered and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent: hexane/DIPE 100/0, 98/2 and 95/5). The pure fractions were collected and the solvent was evaporated [residue; yielding 10.82 g (79%)]. A small amount of the obtained residue was crystallized from DIPE, the precipitate was filtered off and the solvent was evaporated, yielding 1 g of ( $\pm$ )-3,5-dimethylphenyl  $\alpha$ -chlorobenzeneacetate (intermediate 5; mp. 79.0 °C).

#### Example A.5

##### [0066]

a) A mixture of ( $\pm$ )-1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-(1-piperazinyl)piperidine (0.0127 mol), chloroacetonitrile (0.013 mol) and sodium carbonate (0.013 mol) in methylisobutyl keton (110ml) was stirred and refluxed. The mixture was cooled and water was added. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 100/0, 99.5/0.5 and 99/1). The pure fractions were collected and the solvent was evaporated, yielding 3.64g (53%) of ( $\pm$ )-cis-1-[3,5-bis-(trifluoromethyl)benzoyl]-4-[4-(cyanomethyl)-1-piperazinyl]-2-(phenylmethyl)piperidine (intermediate 6).

b) A mixture of intermediate 6 (0.0067 mol) in THF (150 ml) was hydrogenated at 20°C with Raney Nickel (1 g) as a catalyst. After uptake of hydrogen, the catalyst was filtered off and the filtrate was evaporated, yielding 3.77 g of ( $\pm$ )-cis-4-[4-(2-aminoethyl)-1-piperazinyl]-1-[3,5-(trifluoromethyl)benzoyl]-2-(phenylmethyl)piperidine (intermediate 7).

#### Example A.6

[0067] A mixture of 1-(phenylmethyl)-4-piperidinone (0.2 mol) and 1-methylpiperazine (0.2 mol) in methanol (500 ml) was hydrogenated for 8 hours with palladium on activated carbon (10%, 2.5 g) as a catalyst. After uptake of hydrogen, the catalyst was filtered off and the filtrate was evaporated. A mixture of di-*tert*-butyl dicarbonate (0.2 mol) in THF (500 ml) was added to the residue and hydrogenated again with palladium on activated carbon (10%, 2.5 g) as a catalyst. After uptake of hydrogen, the catalyst was filtered off and the filtrate was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 95/5). The pure fractions were collected and the solvent was evaporated, yielding 45.3 g (80%) of 1,1-dimethylethyl 4-(4-methyl-1-piperazinyl)-1-piperidinecarboxylate (intermediate 8).

#### Example A.7

##### [0068]

a) A mixture of 4-methoxypyridine (0.4 mol) in THE (1000 ml) was stirred and cooled in a 2-propanol/CO<sub>2</sub> bath. Ethyl chloroformate (0.4 mol) was added dropwise and the mixture was stirred for 3 hours while cooling (mixture I). In another round-bottom flask, the Grignard-reagent was prepared: Mg (0.44 mol) was stirred in a small amount of (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O. Some 12 was added. A small amount of 1,2-dichloro-4-(chloromethyl)benzene was added. Then, 1,2-dichloro-4-(chloromethyl)benzene (0.4 mol) in (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O (600 ml) was added dropwise at reflux temperature. The mixture was stirred for one hour (mixture II). The Grignard-reagent was decanted off, added to mixture I at < -40 °C, and the resulting reaction mixture was stirred, allowing the temperature to reach RT. The reaction mixture was stirred for one hour at RT. HCl (10 %, 800 ml) was added and the mixture was stirred for 30 minutes, then CH<sub>2</sub>Cl<sub>2</sub> was added. The organic layer was separated, dried, filtered and the solvent evaporated, yielding 57.8 g (44%) of ( $\pm$ )-ethyl 6-[(3,4-dichlorophenyl)methyl]-1,2,3,4-tetrahydro-4-oxo-1-pyridinecarboxylate (intermediate 9).

b) Intermediate 9 (0.176 mol) in THF (880 ml) was stirred under a N<sub>2</sub> flow, and cooled to -78 °C. L-selectride (0.264 mol) was added dropwise at -78 °C. The reaction mixture was stirred for 1 hour, then poured out into water. DIPE was added. The organic layer was separated, washed with an aqueous NaHCO<sub>3</sub> solution, with an aqueous NaCl

solution, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  90/10). The desired fractions were collected and the solvent was evaporated, yielding 20.2 g (34.8%) of ( $\pm$ )-ethyl 2-[(3,4-dichlorophenyl)methyl]-4-oxo-1-piperidinecarboxylate (intermediate 10).

c) Titanium(IV)isopropoxide (0.0269 mol) was added to a mixture of intermediate 10 (0.0224 mol) and intermediate 10 (0.0224 mol) in DCM (11 ml). The mixture was stirred at RT for 3 hours. Sodium cyanoborohydride (0.0224 mol) and then ethanol (10 ml) were added. The mixture was stirred at RT for 48 hours. Water was added and the mixture was stirred.  $\text{CH}_2\text{Cl}_2$  was added and the mixture was stirred. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by HPLC over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  100/0 and 98/2). The pure fractions were collected and the solvent was evaporated. The residue was purified by reversed phase chromatography (eluent:  $\text{NH}_4\text{OAc}$  (0.5% in  $\text{H}_2\text{O}$ )/ $\text{CH}_3\text{OH}$  20/80). Two pure fractions were collected and their solvents were evaporated. The residue was dried and ground, yielding 2 g (16%) of ( $\pm$ )-ethyl trans-2-[(3,4-dichlorophenyl)methyl]-4-[4-[2-[(2,6-dimethylphenyl)amino]-2-oxoethyl]-1-piperazinyl]-1-piperidinecarboxylate (intermediate 11) and 3.5g (28%) of ( $\pm$ )-ethyl cis-2-[(3,4-dichlorophenyl)methyl]-4-[4-[2-[(2,6-dimethylphenyl)amino]-2-oxoethyl]-1-piperazinyl]-1-piperidinecarboxylate (intermediate 12).

d) A mixture of intermediate 11 (0.0034 mol) and potassium hydroxide (0.034 mol) in 2-propanol (150 ml) was stirred and refluxed for 4 days. The solvent was evaporated. The residue was taken up in  $\text{CH}_2\text{Cl}_2$ /water. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$  95/5). The pure fractions were collected and the solvent was evaporated, yielding 0.5 g (30%) of ( $\pm$ )-trans-4-[2-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-N-(2,6-dimethylphenyl)-1-piperazineacetamide (intermediate 13).

#### Example A.8

##### [0069]

a) *Sec*-butyllithium (0.066 mol) was added to a mixture of 1,1-dimethylethyl 1,4-dioxo-8-azaspiro[4.5]-8-carboxylate (0.06 mol) in *N,N,N',N'*-tetramethylethylenediamine (22.6 ml) and  $(\text{C}_2\text{H}_5)_2\text{O}$  (100 ml). The mixture was stirred at  $-70^\circ\text{C}$  for 3 hours. 3,5-difluorobenzaldehyde (0.07 mol) was added dropwise at  $-70^\circ\text{C}$ . The mixture was allowed to warm to RT. Water (50 ml) and DIPE were added. The aqueous layer was separated and extracted with  $\text{CH}_2\text{Cl}_2$ . The combined organic layer was dried, filtered and the solvent was evaporated. Toluene was added and evaporated again, yielding 23 g of ( $\pm$ )-1,1-dimethylethyl 7-[(3,5-difluorophenyl)hydroxymethyl]-1,4-dioxo-8-azaspiro[4.5]-8-carboxylate (intermediate 14)

b) A mixture of intermediate 14 (0.06 mol) and 2-methyl-2-propanol, potassium salt (0.72 g) in toluene (110 ml) was stirred and refluxed for 2 hours. The solvent was evaporated. The residue was stirred in petroleum ether and a small amount of water, and decanted. The residue was dissolved in  $\text{CH}_2\text{Cl}_2$ , dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  100/0, 99/1 and 98/2). Two pure fractions were collected and their solvents were evaporated, yielding 9.2 g (49%) of ( $\pm$ )-3-(3,5-difluorophenyl)tetrahydrospiro[1,3-dioxolan-2,5'(3'H)-1 H-oxazol[3,4-a]pyridin]-1-one (intermediate 15) c)

A mixture of intermediate 15 (0.03 mol) in methanol (250 ml) was hydrogenated at  $50^\circ\text{C}$  with palladium on activated carbon (10%, 2 g) as a catalyst. After uptake of hydrogen, the catalyst was filtered off and the filtrate was evaporated. The residue was purified over silica gel on a glass filter (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  100/0, 98/2 and 95/5 and  $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$  95/5). The desired fractions were collected and the solvent was evaporated, yielding 1.9 g (39 %) of ( $\pm$ )-7-[(3,5-difluorophenyl)methyl]-1,4-dioxo-8-azaspiro[4.5]decane (intermediate 16).

d) A mixture of intermediate 16 (0.012 mol) in  $\text{HCl}$  6N (30ml) was stirred at  $75^\circ\text{C}$  for 2 hours. The mixture was cooled, poured out into ice and a  $\text{NaOH}$  solution and extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was separated, dried and filtered, yielding 2.7 g of ( $\pm$ )-2-[(3,4-difluorophenyl)methyl]-4-piperidinone (intermediate 17). e) A mixture of 3,5-trifluoromethylbenzoyl chloride (0.012 mol) in a small amount of  $\text{CH}_2\text{Cl}_2$  was added dropwise to a stirred mixture of intermediate 17 (0.012 mol) and *N,N*-diethylethanamine (0.024 mol). The mixture was stirred at RT for 1 hour and water was added. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  100/0 and 99.5/0.5). The pure fractions were collected and the solvent was evaporated, yielding 2.7g (48%) of ( $\pm$ )-1-[3,5-bis(trifluoromethyl)benzoyl]-2-[(3,5-difluorophenyl)methyl]-4-piperidinone (intermediate 18).

#### Example A.9

[0070] *Sec*-butyllithium (0.63 mol) was added at  $-78^\circ\text{C}$  under  $\text{N}_2$  flow to a solution of 1,1-dimethylethyl 1,4-dioxo-8-azaspiro[4.5]-8-carboxylate (0.57 mol) and *N,N,N',N'*-tetramethylethylenediamine (1.14 mol) in  $(\text{C}_2\text{H}_5)_2\text{O}$  (1000 ml).



One hour after complete addition, a mixture of 3-(trifluoromethyl)benzaldehyde (0.57 mol) in  $(C_2H_5)_2O$  (200 ml) was added. The mixture was allowed to warm to RT and then stirred at RT for 16 hours. The solvent was evaporated. A mixture of 2-methyl-2-propanol, potassium salt (0.2 mol) in toluene (500 ml) was added. The mixture was stirred at 80°C for 5 hours. The solvent was evaporated. The residue was heated with a saturated  $NH_4Cl$  solution and extracted with  $CH_2Cl_2$ . The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was suspended in DIPE, filtered off and dried. This fraction was dissolved in  $CH_3OH$  (250 ml) and the mixture was hydrogenated with palladium on activated carbon (10%, 3 g) as a catalyst. After uptake of hydrogen, the catalyst was filtered off and the filtrate was evaporated. The residue was purified over silica gel on a glass filter (eluent:  $CH_2Cl_2/CH_3OH$  95/5). The pure fractions were collected and the solvent was evaporated. This fraction was dissolved in HCl (6N, 100 ml) and  $CH_3OH$  (100 ml) and the mixture was stirred at 50°C for 8 hours. The organic solvent was evaporated. The concentrate was washed with a saturated  $K_2CO_3$  solution and extracted with  $CH_2Cl_2$ . The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent:  $CH_2Cl_2/CH_3OH$  95/5). The pure fractions were collected and the solvent was evaporated, yielding 48.5 g (70 %) of  $(\pm)$ -2-[[4-(trifluoromethyl)phenyl]methyl]-4-piperidinone (intermediate 19).

#### Example A.10

[0071]

a) A mixture of ethyl  $\beta$ -oxobenzenebutanoate (0.5 mol) and benzenemethanamine (0.5 mol) in toluene (500 ml) was hydrogenated at 120°C (pressure = 100 kg) overnight in the presence of  $Cu_2Cr_2O_5$  (5 g) and CaO (10 g). After uptake of hydrogen, the catalyst was filtered off and the filtrate was evaporated, yielding 29.7 g of  $(\pm)$ -ethyl N,2-bis(phenylmethyl)- $\beta$ -alanine (intermediate 20).

b) Ethyl chloroacetate (0.3 mol) was added to a mixture of intermediate 20 (0.2 mol) in DMF (250 ml). The mixture was stirred and triethylamine (0.4 mol) was added. The mixture was stirred at 60°C overnight. The solvent was evaporated and the residue was taken up in water/ $CH_2Cl_2$ . The organic layer was separated, dried, filtered and the solvent was evaporated, yielding 76.6 g of  $(\pm)$ -ethyl 3-[(2-ethoxy-2-oxoethyl)(phenylmethyl)amino]benzenebutanoate (intermediate 21).

c) Intermediate 21 (0.2 mol) was heated to 80°C under  $N_2$  flow.  $NaOCH_3$  (44 g) was added. The mixture was stirred at 80°C for 30 minutes. The solvent was evaporated and water (170 ml) and HCl (6N, 60 ml) were added. The mixture was stirred and refluxed for 1 hour, then cooled, alkalized with NaOH and extracted with  $CH_2Cl_2$ . The organic layer was separated, washed with water and a saturated NaCl solution, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent:  $CH_2Cl_2/CH_3CN$  100/0 to 96/4). The pure fractions were collected and the solvent was evaporated, yielding 7.8 g of  $(\pm)$ -1,5-bis(phenylmethyl)-3-pyrrolidinone (intermediate 22).

d) A mixture of intermediate 22 (0.027 mol) and  $CH_3SO_3H$  (0.03 mol) in THF (200 ml) was hydrogenated with palladium on activated carbon (10%, 2 g) as a catalyst. After uptake of hydrogen, the catalyst was filtered off, yielding  $(\pm)$ -5-(phenylmethyl)-3-pyrrolidinone methanesulfonate(1:1) (intermediate 23).

e) 3,5-di(trifluoromethyl)benzoyl chloride (0.03 mol) was added to intermediate 23 (0.027 mol). The mixture was stirred and triethylamine (0.1 mol) was added. The mixture was stirred at RT for 18 hours and then washed with water, NaOH and a saturated NaCl solution. The organic layer was separated, washed with a saturated NaCl solution, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent:  $CH_2Cl_2/CH_3OH$  98/2). The pure fractions were collected and the solvent was evaporated, yielding 1.4 g of  $(\pm)$ -1-[3,5-bis(trifluoromethyl)benzoyl]-5-(phenylmethyl)-3-pyrrolidinone (intermediate 24).

#### B. Preparation of the compounds of formula (I)

##### Example B.1

[0072]

a) Titanium(IV)isopropoxide (16.5 g) was added to a mixture of intermediate 3 (21.5 g) and 1-(phenylmethyl)piperazine (8.81 g) in DCM (35 ml). The mixture was stirred for 3 hours at RT. Sodium cyanoborohydride (2.85 g) and ethanol (70 ml) were added and the resulting reaction mixture was stirred overnight at RT. Water (5 ml) and DCM were added. The biphasic mixture was filtered over dicalite, and the filter residue was washed with DCM. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was crystallized from  $CH_3CN$  and the precipitate was filtered off and dried, yielding 7.93 g (26.9 %) of  $(\pm)$ -cis-1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-[4-(phenylmethyl)-1-piperazinyl]piperidine (compound 16; mp. 143.8 °C).

b) The mother liquor was concentrated and the residue was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  100/0, then 99/1, 98/2, 97/3). The desired fractions ((A) and (B)) were collected and their solvent was evaporated. The A-isomer was crystallized from  $\text{CH}_3\text{CN}$ , filtered off and dried, yielding 1.11 g (4 %) of compound 16. The pure fractions of the B-isomer were concentrated, yielding 5.9 g (20%) of ( $\pm$ )-*trans*-1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-[4-(phenylmethyl)-1-piperazinyl]piperidine. The impure fractions of the B-isomer were collected and the solvent was evaporated. The residue was converted into the fumaric acid salt (1:2) in ethanol. The precipitate was filtered off and dried, yielding 1.89 g ( $\pm$ )-*trans*-1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-[4-(phenylmethyl)-1-piperazinyl]piperidine (E)-2-butenedioate(1:2) (compound 17; mp. 240.3 °C).

#### Example B.2

[0073] A mixture of compound 16 (8.4 g) in methanol (250 ml) was hydrogenated at 50 °C with palladium on activated carbon (10 %) (2 g) as a catalyst. After uptake of  $\text{H}_2$ , the catalyst was filtered off and the filtrate was evaporated, yielding 7 g (100 %) of ( $\pm$ )-1-[3,5-bis-(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-(1-piperazinyl)piperidine (compound 15).

#### Example B.3

[0074]

a) Titanium(IV)isopropoxide (13.2 g) was added to a mixture of intermediate 3 (17.16 g) and *N*-(2,6-dimethylphenyl)-1-piperazineacetamide (9.88 g) in DCM (20 ml). This mixture was stirred for 3 hours at RT. Sodium cyanoborohydride (2.52 g) in ethanol (20 ml) was added and the resulting reaction mixture was stirred overnight at RT. Water (10 ml) was added and the reaction mixture was extracted with DCM (800 ml). The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was taken up into water and this mixture was extracted with DCM. The separated organic layer was dried, filtered, and the solvent evaporated. The residue was pre-purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  97/3). The desired fractions were collected and the solvent was evaporated, giving 4 g of the *trans*-racemate. Resolution was obtained by purification over stationary phase Chiralcel OD (eluent:  $\text{CH}_3\text{OH}$  100%). Two desired *trans*-fraction groups were collected and their solvent was evaporated, yielding 1.75 g fraction 1 and 2 g fraction 2. Fraction 1 was dissolved in DCM, filtered and the filtrate was evaporated. The residue was dried, yielding 1.55 g (6%) (-)-(A)-*trans*-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]-*N*-(2,6-dimethylphenyl)-1-piperazine acetamide (compound 26; mp. 97.4 °C;  $[\alpha]_{\text{D}}^{20} = -5.81^\circ$  (c = 1 % in DMF)). Fraction 2 was dissolved in DCM, filtered and the filtrate was evaporated. The residue was dried, yielding 1.70 g (6%) (+)-(B)-*trans*-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]-*N*-(2,6-dimethylphenyl)-1-piperazine acetamide (compound 27; mp. 96.8 °C;  $[\alpha]_{\text{D}}^{20} = +5.71^\circ$  (c = 1 % in DMF)).

b) Compound 27 was dissolved in warm 2-propanol and converted into the (L)-malic acid salt with a solution of (L) malic acid in 2 propanol. The mixture was stirred for 2 hours and the precipitate was filtered off and dried, yielding (+)-(B)-*trans*-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]-*N*-(2,6-dimethylphenyl)-1-piperazine acetamide (L)-malic acid (1:1) (compound 95).

#### Example B.4

[0075] A mixture of compound 15 (2.5 g), intermediate 5 (1.65 g) and sodium carbonate (0.64 g) in methylisobutylketone (50 ml) was stirred and refluxed for 3 hours. The reaction mixture was washed and the separated organic layer was dried, filtered and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  100/0 and 99.5/0.5). The pure fractions were collected and the solvent was evaporated, yielding 1.59 g (43%) of ( $\pm$ )-3,5-dimethylphenyl *cis*-4-[1-[3,5-bis-(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]- $\alpha$ -phenyl-1-piperazineacetate (compound 43; mp. 88.1 °C).

#### Example B.5

[0076] A mixture of intermediate 2 (3.2 g), 1-(diphenylmethyl)piperazine (2.5 g) and aluminum tributoxide (2 g) in toluene (250 ml) was hydrogenated for 48 hours at 50 °C, with palladium on activated carbon (10 %; 2 g) as a catalyst in the presence of thiophene (4 % solution; 1 ml). After uptake of hydrogen (1 equiv), the catalyst was filtered off and the filtrate was evaporated. The residue was purified by high-performance liquid chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$  100/0, upgrading to 90/10). Two pure fractions were collected and their solvent was evaporated, resulting in residue 1 and residue 2. Residue 1 was suspended in DIPE. The precipitate was filtered off and dried, yielding

0.94 g (17%) of ( $\pm$ )-*cis*-1-(dimethylbenzoyl)-4-[4-(diphenylmethyl)-1-piperazinyl]-2-(phenylmethyl)piperidine (compound 12; mp. 100.8°C). Residue 2 was dried, yielding 0.2 g (3.6%) of ( $\pm$ )-*trans*-1-(dimethylbenzoyl)-4-[4-(diphenylmethyl)-1-piperazinyl]-2-(phenylmethyl)piperidine (compound 13).

#### Example B.6

[0077] A mixture of compound 15 (0.005 mol) and 1,2-epoxyethylbenzene (0.006 mol) in methanol (50 ml) was stirred at RT for 1 hour. The mixture was stirred and refluxed for 3 hours. The solvent was evaporated and the residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 100/0, 99/1 and 98/2). The pure fractions were collected and the solvent was evaporated. The residue was purified by HPLC over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 98/2 to 95/5). Two pure fractions were collected and their solvents were evaporated. Each residue was dried, yielding 0.7 g (23%) of ( $\pm$ )-*cis*-1-[3,5-bis(trifluoromethyl)benzoyl]-4-[4-(2-hydroxy-2-phenylethyl)-1-piperazinyl]-2-(phenylmethyl)piperidine (compound 60) and 0.23 g (7%) of ( $\pm$ )-*cis*-1-[3,5-bis(trifluoromethyl)benzoyl]-4-[4-(2-hydroxy-1-phenylethyl)-1-piperazinyl]-2-(phenylmethyl)piperidine (compound 61).

#### Example B.7

[0078] Compound 15 (0.005 mol), 2-chloro-1-[(2-methyl-5-oxazolyl)methyl]-1H-benzimidazole (0.005 mol) and Copper (0.005 mol) were stirred at 140°C for 2 hours. The mixture was cooled, dissolved in CH<sub>2</sub>Cl<sub>2</sub>, filtered and washed with CH<sub>2</sub>Cl<sub>2</sub> and a diluted NH<sub>4</sub>OH solution. The organic layer was separated, washed with a diluted NH<sub>4</sub>OH solution, dried, filtered and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 100/0, 99.5/0.5, 99/1, 98.5/1.5 and 98/2). The pure fractions were collected and the solvent was evaporated. The residue was dried, yielding 1.42 g (40%) ( $\pm$ )-*cis*-1-[3,5-bis(trifluoromethyl)benzoyl]-4-[4-[1-[(2-methyl-5-oxazolyl)methyl]-1H-benzimidazol-2-yl]-1-piperazinyl]-2-(phenylmethyl)piperidine (compound 70).

#### Example B.8

[0079] A mixture of intermediate 7 (0.0033 mol) and 3,5-dimethylbenzoyl chloride (0.0035 mol) in DCM (50 ml) was stirred at RT for 15 minutes. Triethylamine (0.007 mol) was added and the mixture was stirred at RT for 1 hour. Water was added. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was converted into the fumaric acid salt (1:1) with 2-propanol. The precipitate was filtered off and dried. The residue was converted into the free base with NaOH. The precipitate was filtered off and dried. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 100/0, 99.5/0.5, 99/1, 98/2 and 97/3). The pure fractions were collected and the solvent was evaporated. The residue was dried, yielding 0.8 g (36%) ( $\pm$ )-*cis*-N-[2-[4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]-1-piperazinyl]ethyl]-3,5-dimethylbenzamide (compound 116).

#### Example B.9

[0080] A mixture of compound 74, prepared according to example B.4, (0.004 mol) in methanol (150 ml) was hydrogenated at 50°C with palladium on activated carbon (10 %; 1 g) as a catalyst in the presence of thiophene (4 % solution, 1 ml). After uptake of hydrogen, the catalyst was filtered off and the filtrate was evaporated. The residue was crystallized from DIPE. The precipitate was filtered off, washed with DIPE and dried. This fraction was dissolved in toluene. The mixture was filtered and the solvent was evaporated. The residue was suspended in DIPE. The precipitate was filtered off and dried. This fraction was converted into the fumaric acid salt (1:2) with a warm solution of fumaric acid (0.52g) in ethanol. The mixture was stirred for 6 hours. The precipitate was filtered off and dried, yielding 0.91 g (25%) of ( $\pm$ )-*cis*-N-(4-amino-2,6-dimethylphenyl)-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]-1-piperazineacetamide (E)-2-butenedioate(1:2) (compound 129).

#### Example B.10

[0081] *Sec*-butyllithium (0.055 mol) was added at -78°C under N<sub>2</sub> flow to a solution of 1,1-dimethylethyl 4-(4-methyl-1-piperazinyl)-1-piperidinecarboxylate (0.05 mol) and *N,N,N',N'*-tetramethylethylenediamine (0.1 mol) in (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O (50 ml). 2 hours after complete addition, a mixture of benzaldehyde (0.05 mol) in (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O (50 ml) was added. The mixture was allowed to warm to RT and then stirred at 25°C for 16 hours. The solvent was evaporated and the residue was washed with a saturated NH<sub>4</sub>Cl solution and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried, filtered and the solvent was evaporated. A solution of 2-methyl-2-propanol, potassium salt (0.02 mol) in toluene (100 ml) was added to this fraction and the mixture was stirred at 100°C for 2 hours. The solvent was evaporated. The residue was washed with a saturated NH<sub>4</sub>Cl solution, extracted with CH<sub>2</sub>Cl<sub>2</sub> and decanted. The organic layer was dried, filtered

and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 95/5). The pure fractions were collected and the solvent was evaporated. This fraction was dissolved in methanol (150 ml) and hydrogenated with palladium on activated carbon (10%, 3g) as a catalyst. After uptake of hydrogen, the catalyst was filtered off and the filtrate was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 95/5). The pure fractions were collected and the solvent was evaporated. This fraction was dissolved in DCM (20 ml) and Triethylamine (2 ml). 3,5-di(trifluoromethyl)benzoyl chloride (0.0087 mol) was added at 0°C. 1 hour after complete addition, water was added and the mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 95/5). The pure fractions were collected and the solvent was evaporated. This fraction was converted into the (E)-2-butenedioic acid salt (1:2) with ethanol. The precipitate was filtered off and dried, yielding 4.7 g (74%) of (±)-*cis*-1-[3,5-bis(trifluoromethyl)benzoyl]-4-(4-methyl-1-piperazinyl)-2-(phenylmethyl)piperidine (E)-2-butenedioate(1:2) (compound 130).

#### Example B.11

**[0082]** A mixture of compound 15 (0.005 mol), *N*-[2-(3,4-dichlorophenyl)-4-[(methylsulfonyl)oxy]butyl]-*N*-methyl benzamide (0.0055 mol) and NaHCO<sub>3</sub> (0.0055 mol) in ethanol (50 ml) as stirred and refluxed for 6 hours. The solvent was evaporated, the residue was taken up in water and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 100/0, 99/1, 98/2 and 97/3). The pure fractions were collected and the solvent was evaporated. The residue was converted into the fumaric acid salt (1:2) with ethanol. The precipitate was filtered off and dried, yielding 1.42 g (27%) of (±)-*cis*-*N*-[4-[4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]-1-piperazinyl]-2-(3,4-dichlorophenyl)butyl]-*N*-methylbenzamide (E)-2-butenedioate(1:2) (compound 93).

#### Example B.12

**[0083]** A mixture of (±)-1-[3,5-bis(trifluoromethyl)benzoyl]-2-[(3,5-difluorophenyl)methyl]-4-piperidinone (0.0058 mol), *N*-(2,6-dimethylphenyl)-1-piperazineacetamide (0.0058 mol) and titanium(IV)isopropoxide (0.0064 mol) in 2-propanol (5 ml) was stirred at RT overnight. NaBH<sub>4</sub> (0.0116 mol) and ethanol (15 ml) were added. The mixture was stirred for 2 days. Water (5 ml) was added and the mixture was stirred for 10 minutes. CH<sub>2</sub>Cl<sub>2</sub> (200 ml) was added. The organic layer was separated, dried, filtered and the solvent was evaporated. This fraction was purified by HPLC over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 98/2 to 90/10 over a 30-minute period). Two pure fractions (F1 and F2) were collected and their solvents were evaporated. F1 was purified by column chromatography over RP18 (eluent: NH<sub>4</sub>OAc (0.5% in H<sub>2</sub>O)/CH<sub>3</sub>CN 40/60). The pure fractions were collected and the solvent was evaporated. The residue was dried, yielding 0.33 g (8%) of (±)-*cis*-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-[(3,5-difluorophenyl)methyl]-4-piperidinyl]-*N*-(2,6-dimethylphenyl)-1-piperazineacetamide (compound 132). F2 was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 100/0 to 92/8 over a 30-minute period). The pure fractions were collected and the solvent was evaporated. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub>, filtered and the solvent was evaporated. The residue was dried, yielding 0.24 g (6%) of (±)-*trans*-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-[(3,5-difluorophenyl)methyl]-4-piperidinyl]-*N*-(2,6-dimethylphenyl)-1-piperazineacetamide (compound 133).

#### Example B.13

**[0084]** 3,5-di(trifluoromethyl)benzoyl chloride (0.0011 mol) was added to a mixture of (±)-*trans*-4-[2-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-*N*-(2,6-dimethylphenyl)-1-piperazineacetamide (0.001 mol) in DCM (20ml). The mixture was stirred for 5 minutes. Triethylamine (2ml) was added. The mixture was stirred at RT for 3 hours, washed with a diluted NaOH solution and with water, and then dried. The solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 96/4). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from CH<sub>3</sub>CN. The precipitate was filtered off and dried, yielding 0.32 g (44%) of (±)-*trans*-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-*N*-(2,6-dimethylphenyl)-1-piperazineacetamide (compound 139).

#### Example B.14

**[0085]** A mixture of compound 15 (0.01 mol) and imidazo[1,2-*a*]pyridin-2-carboxaldehyde (0.01 mol) in methanol (250 ml) was hydrogenated at RT overnight with palladium on activated carbon (10%, 2 g) as a catalyst in the presence of thiophene (4% solution, 2 ml). After uptake of hydrogen, the catalyst was filtered off and the filtrate was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 110/0, 99/1, 98/2, 97/3 and 96/4). The

pure fractions were collected and the solvent was evaporated. The residue was converted into the fumaric acid salt (1:2) from ethanol. The precipitate was filtered off and dried, yielding 2.8 g (32%) of ( $\pm$ )-*cis*-1-[3,5-bis(trifluoromethyl)benzoyl]-4-[4-(imidazo[1,2-a]pyridin-2-ylmethyl)-1-piperazinyl]-2-(phenylmethyl)piperidine (E)-2-butenedioate(1:2) (compound 111).

#### Example B.15

[0086] (+)-(B-*trans*)-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]-*N*-(2,6-dimethylphenyl)-1-piperazineacetamide (0.003 mol) was dissolved in ethanol (20 ml). A solution of fumaric acid (0.003 mol) in ethanol (15 ml) was added and the mixture was stood for 7 days. The precipitate was filtered off and dried, yielding 1.2 g of (B-*trans*)-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]-*N*-(2,6-dimethylphenyl)-1-piperazineacetamide (E)-2-butenedioate(1:1) (compound 128).

#### Example B.16

[0087] A mixture of ( $\pm$ )-1-[3,5-bis(trifluoromethyl)benzoyl]-5-(phenylmethyl)-3-pyrrolidinone (0.0037 mol) and *N*-(2,6-dimethylphenyl)-1-piperazineacetamide (0.0037 mol) in methanol (150 ml) was hydrogenated at 50°C with palladium on activated carbon (10%, 1 g) as a catalyst in the presence of thiophene solution (1 ml). After uptake of hydrogen, the catalyst was filtered off and the filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 95/5). The desired fractions were collected and the solvent was evaporated. The residue was dried and then crystallized from DIPE. The precipitate was filtered off and dried, yielding 0.35 g (15 %) of ( $\pm$ )-*cis*-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-5-(phenylmethyl)-3-pyrrolidinyl]-*N*-(2,6-dimethylphenyl)-1-piperazineacetamide (compound 131).

#### Example B.17

[0088] ( $\pm$ )-*cis*-1-(phenylmethyl)-4-[2-(phenylmethyl)-1-piperidinyl]piperazine (0.00043 mol) was added to 3,4-dichlorobenzeneacetic acid ( $\pm$  0.0004 mol) and 1-hydroxybenzotriazole hydrate (0.080 g) in DCM (5 ml). The mixture was stirred and cooled on an ice/ethanol-bath, under N<sub>2</sub> flow. Triethylamine was added dropwise. A solution of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.120 g) in DCM (5 ml) was added and the reaction mixture was allowed to warm to RT, under N<sub>2</sub>. The reaction mixture was stirred overnight. The mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub>, until a 15-ml total volume was obtained. Then, the compound was isolated and purified by HPLC over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub> to CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 90/10 over 20 minutes at 125 ml/minute). The desired fractions were collected and the solvent was evaporated, yielding 0.020 g of ( $\pm$ )-*cis*-1-[(3,4-dichlorophenyl)acetyl]-2-(phenylmethyl)-4-[4-(phenylmethyl)-1-piperazinyl]piperidine (compound 181).

#### Example B.18


[0089] 3,5-di(trifluoromethyl)-1-isocyanatobenzene (0.0025 mol) in DCM (10 ml) was added to a mixture of ( $\pm$ )-*trans*-*N*-(2,6-dimethylphenyl)-4-[2-(phenylmethyl)-4-piperidinyl]-1-piperazineacetamide (0.0025 mol) in DCM (15 ml). The mixture was stirred at RT overnight. The precipitate was filtered off and dried, yielding 0.66 g (40%) of ( $\pm$ )-*trans*-4-[1-[[[3,5-bis(trifluoromethyl)phenyl]amino]carbonyl]-2-(phenylmethyl)-4-piperidinyl]-*N*-(2,6-dimethylphenyl)-1-piperazineacetamide (compound 143).

#### Example B.19

[0090] A mixture of ( $\pm$ )-1-[3,5-bis(trifluoromethyl)benzoyl]-2-[[3-fluoro-5-(trifluoromethyl)phenyl]methyl]-4-piperidinone (0.01 mol) and *N*-(2,6-dimethylphenyl)-1-piperazineacetamide (0.01 mol) in 2-propanol (150 ml) was hydrogenated at 50°C with platinum on activated carbon (5 %, 2 g) as a catalyst in the presence of titanium(IV)isopropoxide (2.84 g) and thiophene solution (1 ml). After uptake of hydrogen, the catalyst was filtered off and the filtrate was evaporated. The residue was taken up in CH<sub>2</sub>Cl<sub>2</sub> and H<sub>2</sub>O. The organic layer was separated, washed several times with H<sub>2</sub>O, dried, filtered over dicalite and the solvent was evaporated. This fraction was purified by HPLC over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 98/2). Two pure fractions were collected and their solvents were evaporated. The residue was dried, yielding 0.72 g (10 %) of ( $\pm$ )-*cis*-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-[[3-fluoro-5-(trifluoromethyl)phenyl]methyl]-4-piperidinyl]-*N*-(2,6-dimethylphenyl)-1-piperazineacetamide (compound 140) and 0.88 g (12 %) of ( $\pm$ )-*trans*-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-[[3-fluoro-5-(trifluoromethyl)phenyl]methyl]-4-piperidinyl]-*N*-(2,6-dimethylphenyl)-1-piperazineacetamide (compound 141).

[0091] Tables 1 to 4 list compounds of formula (I) that were prepared according to one or more of the foregoing

examples (Ex.).

Table 1

Co. No.	Ex.	-L	Physical data (mp = melting point)
1	6	-H	(±)- <i>cis</i>
2	6	-H	(±)- <i>trans</i>
3	9		(±)- <i>cis</i> ; mp 196.9 °C
4	9		(±)- <i>trans</i>
5	5		(±)- <i>cis</i> ; mp 94.0 °C
6	5		(±)- <i>trans</i>
7	8		(±)- <i>cis</i> -(E) ; mp 201.0 °C





















Co. No.	Ex.	-L	Physical data (mp = melting point)
8	8		(±)- <i>trans</i> -(E); mp 210.1 °C
9	8		(±)- <i>cis</i> ; mp 92.1 °C
10	9		(±)- <i>cis</i> ; mp 72.8 °C
11	9		(±)- <i>trans</i>
12	9		(±)- <i>cis</i> ; mp 100.8 °C
13	9		(±)- <i>trans</i>













Table 2
























Co. No.	Ex.	n	p	-L	Physical data (mp = melting point)
14	B.2	1	1	-H	(±)- <i>trans</i>
15	B.2	1	1	-H	(±)- <i>cis</i>
16	B.1.a	1	1		(±)- <i>cis</i> ; mp 143.8 °C
17	B.1.a+b	1	1		(±)- <i>trans</i> ; mp 240.3 °C; fumaric acid (1:2)











Co. No.	Ex.	n	p	-L	Physical data (mp = melting point)
18	B.1	1	1		(±)- <i>cis</i> ; mp 120°C
19	B.1	1	1		(±)- <i>trans</i> ; mp 150°C
20	B.1	1	1		(±)- <i>cis</i> ; mp 70.4°C
21	B.1	1	1		(±)- <i>trans</i> ; mp 169.1°C
22	B.1	1	1		(±)- <i>trans</i> ; mp 173.8°C
23	B.1	1	1		(±)- <i>cis</i> ; mp 93.2°C
24	B.1	1	1		(±)- <i>trans</i> ; mp 100.1°C
25	B.1	1	1		(±)- <i>trans</i> ; mp 75.4°C
26	B.1 and B.3	1	1		(-)-(A)- <i>trans</i> ; mp 97.4°C; [α] <sub>D</sub> <sup>20</sup> = -5.81° (c = 1 % in DMF)
27	B.1 and B.3	1	1		(+)-(B)- <i>trans</i> ; mp 96.8°C; [α] <sub>D</sub> <sup>20</sup> = +5.71° (c = 1 % in DMF)
28	B.1	1	1		(±)- <i>cis</i> ; (E)













Co. No.	Ex.	n	p	-L	Physical data (mp = melting point)
29	B.1	1	1		(±)- <i>trans</i> ; (E)
30	B.1	1	1		(±)- <i>trans</i> ; mp 185.7°C
31	B.1	1	1		(±)- <i>cis</i> ; mp 77.5°C
32	B.1	1	1		(±)- <i>cis</i> ; mp 183.1°C
33	B.1	1	1		(±)- <i>trans</i> ; mp 115.6°C
34	B.1	1	2		(±)- <i>trans</i> ; mp 120.1°C
35	B.1	1	1		(±)- <i>cis</i> ; mp 150.9°C
36	B.1	1	1		(±)- <i>trans</i> ; mp 120.8°C
37	B.4	1	1		(±)- <i>cis</i> ; mp 85.6°C
38	B.4	1	1		(±)- <i>trans</i> ; mp 170.5°C
39	B.4	1	1		(±)- <i>cis</i> ; mp 192.9°C
40	B.4	1	1		(±)- <i>trans</i> ; mp 240.7°C










Co. No.	Ex.	n	p	-L	Physical data (mp = melting point)
41	B.4	1	1		(+)-(A)- <i>cis</i> ; mp 177.3°C; [α] <sub>D</sub> <sup>20</sup> = +19.88° (c = 1 % in methanol)
42	B.4	1	1		(-)-(B)- <i>cis</i> ; mp 177.3°C; [α] <sub>D</sub> <sup>20</sup> = -20.34° (c = 1 % in methanol)
43	B.4	1	1		(±)- <i>cis</i> ; mp 88.1°C
44	B.4	1	1		(±)- <i>cis</i> ; mp 227.1°C; fumaric acid (1:2)
45	B.4	1	1		(±)- <i>trans</i> ; mp 200.2°C
46	B.4	1	1		(±)- <i>trans</i> ; mp 105.6°C
47	B.4	1	1		(±)- <i>cis</i> ; mp 89.2°C
48	B.1	1	1		(±)- <i>trans</i> ; mp 89.7°C
49	B.1	1	1		(±)- <i>cis</i> ; mp 135.8°C
50	B.4	1	1		(±)- <i>trans</i> ; mp 140.4°C

Co. No.	Ex.	n	p	-L	Physical data (mp = melting point)
51	B.4	1	1		(±)- <i>cis</i> ; mp 173.5°C
52	B.4	1	1		(±)- <i>cis</i> ; mp 101.5°C
53	B.4	1	1		(±)- <i>trans</i> ; mp 185.8°C
54	B.5	1	1		(±)- <i>cis</i> ; mp 260°C
55	B.5	1	1		(±)- <i>trans</i> ; mp 75.2°C
56	B.5	1	1		(±)- <i>trans</i> ; mp 80.1°C
57	B.5	1	1		(±)- <i>cis</i>
58	B.1	1	1		(±)
59	B.4	1	1		(±)- <i>cis</i> ; mp 106.4°C
60	B.6	1	1		(±)- <i>cis</i>
61	B.6	1	1		(±)- <i>cis</i>

Co. No.	Ex.	n	p	-L	Physical data (mp = melting point)
62	B.1	1	1	$\text{CH}_2\text{-CH}_2\text{-O-CH}_2\text{CH}_3$ 	(±)- <i>cis</i>
63	B.1	1	2		(±)- <i>cis</i> ; fumaric acid (1:2)
64	B.2	1	2	-H	(±)- <i>cis</i>
65	B.4	1	2		(±)- <i>cis</i>
66	B.1	1	1		(±)- <i>cis</i>
67	B.1	1	1		(±)- <i>trans</i>
68	B.1	1	2		(±)- <i>trans</i> ; fumaric acid (1:2)
69	B.4	1	1		(±)- <i>trans</i>
70	B.7	1	1		(±)- <i>cis</i>
71	B.4	1	1		(±)- <i>cis</i>
72	B.4	1	1		(±)- <i>cis</i>

Co. No.	Ex.	n	p	-L	Physical data (mp = melting point)
73	B.4	1	1		(±)-cis; fumaric acid (1:2)
74	B.4	1	1		(±)-cis
75	B.4	1	1		(±)-cis; fumaric acid (1:1)
76	B.4	1	1	$-\text{CH}_2-\text{C}(=\text{O})-\text{NH}-\text{CH}(\text{CH}_3)_2$	(±)-cis
77	B.2	1	2	-H	(±)-trans
78	B.4	1	2		(±)-trans
79	B.8	1	1		(±)-cis
80	B.1	1	1		(+)-(B)-trans
81	B.2	1	1	-H	(+)-(B)-trans
82	B.1	1	1		(±)-cis
83	B.1	1	1		(±)-trans
84	B.4	1	1		(±)-cis

Co. No.	Ex.	n	p	-L	Physical data (mp = melting point)
85	B.7	1	1		(±)- <i>trans</i>
86	B.6	1	1		(±)- <i>trans</i>
87	B.1	1	1		(±)- <i>trans</i>
88	B.1	1	1		(±)- <i>cis</i> ; fumaric acid (1:2)
89	B.4	1	1		(±)- <i>trans</i>
90	B.11	1	1		(±)- <i>trans</i> ; fumaric acid (1:2)
91	B.4	1	1		(±)- <i>trans</i> ; fumaric acid (1:2)
92	B.11	1	1		(±)- <i>trans</i>
93	B.11	1	1		(±)- <i>cis</i> ; fumaric acid (1:2)
94	B.4	1	1		(±)- <i>trans</i>

Co. No.	Ex.	n	p	-L	Physical data (mp = melting point)
95		1	1		(B)- <i>trans</i> ; (L)-malic acid (1:1)
96	B.4	1	1		(±)- <i>trans</i> ; fumaric acid (1:2)
97	B.4	1	1		(±)- <i>trans</i>
98	B.4	1	1		(±)- <i>cis</i>
99	B.4	1	1		(±)- <i>cis</i>
100	B.4	1	1		(±)- <i>cis</i>
101	B.4	1	1		(±)- <i>cis</i>
102	B.4	1	1		(±)- <i>cis</i>
103	B.4	1	1		(±)- <i>cis</i> fumaric acid (1:2)


Co. No.	Ex.	n	p	-L	Physical data (mp = melting point)
104	B.4	1	1		(±)- <i>trans</i>
105	B.4	1	1		(±)- <i>trans</i>
106	B.4	1	1		(±)- <i>trans</i>
107	B.4	1	1		(±)- <i>trans</i>
108	B.4	1	1		(±)- <i>trans</i>
109	B.4	1	1		(±)- <i>trans</i>
110	B.14	1	1		(±)- <i>trans</i> ; fumaric acid (1:2)
111	B.14	1	1		(±)- <i>cis</i> ; fumaric acid (1:2)
112	B.4	1	1		(±)- <i>trans</i>
113	B.1	1	1		(±)- <i>trans</i>
114	B.2	1	1	H	(±)- <i>trans</i> ; fumaric acid (1:2)
115	B.8	1	1		(±)- <i>trans</i>





Co. No.	Ex.	n	p	-L	Physical data (mp = melting point)
116	B.8	1	1		(±)- <i>cis</i>
117	B.8	1	1		(±)- <i>trans</i>
118	B.4	1	1		(±)- <i>trans</i> ; fumaric acid (1:2)
119	B.4	1	1		(±)- <i>trans</i>
120	B.13	1	1		(±)- <i>trans</i>
121	B.1	1	1		(±)- <i>cis</i>
122	B.1	1	1		(±)- <i>trans</i>
123	B.4	1	1		(±)- <i>cis</i>
124	B.15	1	1		(B)- <i>trans</i> ; benzoate (1:1)

Co. No.	Ex.	n	p	-L	Physical data (mp = melting point)
125	B.15	1	1		(B)- <i>trans</i> ; maleic acid (1:1)
126	B.15	1	1		(B)- <i>trans</i> ; hydrochloric acid (1:2) hydrate (1:1)
127	B.15	1	1		(B)- <i>trans</i> ; succinic acid (1:1)
128	B.15	1	1		(B)- <i>trans</i> ; fumaric acid (1:1)
129	B.9	1	1		(±)- <i>cis</i> ; fumaric acid (1:2)
130	B.10	1	1	-CH <sub>3</sub>	(±)- <i>cis</i> ; fumaric acid (1:2)
131	B.16	0	1		(±)- <i>cis</i>

Table 3







Co. No.	Ex.	X&	R <sup>2</sup>	R <sub>a</sub>	R <sub>b</sub>	R <sub>c</sub>	Physical data
132	B.12	c.b.	3,5-di(trifluoro- methyl)phenyl	F	H	F	(±)- <i>cis</i>
133	B.12	c.b.	3,5-di(trifluoro- methyl)phenyl	F	H	F	(±)- <i>trans</i>
134	B.12	c.b.	3,5-di(trifluoro- methyl)phenyl	H	F	F	(±)- <i>cis</i>
135	B.12	c.b.	3,5-di(trifluoro- methyl)phenyl	H	F	F	(±)- <i>trans</i>
136	B.4	c.b.	3,5-di(trifluoro- methyl)phenyl	H	CF <sub>3</sub>	H	(±)-(B) fumaric acid (1:1)
137	B.4	c.b.	3,5-di(trifluoro- methyl)phenyl	H	CF <sub>3</sub>	H	(±)-(A)
138	B.13	c.b.	3,5-di(trifluoro- methyl)phenyl	H	Cl	Cl	(±)- <i>cis</i>
139	B.13	c.b.	3,5-di(trifluoro- methyl)phenyl	H	Cl	Cl	(±)- <i>trans</i>
140	B.19	c.b.	3,5-di(trifluoro- methyl)phenyl	F	H	CF <sub>3</sub>	(±)- <i>cis</i>
141	B.19	c.b.	3,5-di(trifluoro- methyl)phenyl	F	H	CF <sub>3</sub>	(±)- <i>trans</i>
142	B.13	c.b.	3-isopropoxyphenyl	H	H	H	(±)- <i>cis</i>
143	B.18	-NH-	3,5-di(trifluoro- methyl)phenyl	H	H	H	(±)- <i>trans</i>
144	B.13	c.b.	phenyl	H	H	H	(±)- <i>cis</i>
145	B.13	c.b.	2-naphthyl	H	H	H	(±)- <i>trans</i>
146	B.13	c.b.	2-quinoliny	H	H	H	(±)- <i>trans</i>
147	B.13	c.b.	2-quinoxaliny	H	H	H	(±)- <i>trans</i>
148	B.13	-O-	benzyl	H	H	H	(±)- <i>trans</i>
149	B.17	c.b.	3-methyl- benzofuran-2-yl	H	H	H	(±)- <i>trans</i>
150	B.17	c.b.	5-fluoro-indol-2-yl	H	H	H	(±)- <i>trans</i>
151	B.17	c.b.	5-indolyl	H	H	H	(±)- <i>trans</i>
152	B.17	c.b.	5-methyl-pyrazin-2- yl	H	H	H	(±)- <i>trans</i>
153	B.13	c.b.	phenyl	H	H	H	(±)- <i>trans</i>

Co. No.	Ex.	X &	R <sup>2</sup>	R <sub>a</sub>	R <sub>b</sub>	R <sub>c</sub>	Physical data
154	B.13	c.b.	5-methyl-isoxazol-3-yl	H	H	H	(±)- <i>trans</i>
155	B.13	c.b.	2,4,6-trimethylphenyl	H	H	H	(±)- <i>cis</i>
156	B.13	c.b.	3,4,5-trimethoxyphenyl	H	H	H	(±)- <i>cis</i>
157	B.13	c.b.	3-cyanophenyl	H	H	H	(±)- <i>cis</i>
158	B.13	c.b.		H	H	H	(±)- <i>cis</i>
159	B.13	c.b.	3,5-difluorophenyl	H	H	H	(±)- <i>cis</i>
160	B.13	c.b.	2,6-dichloro-pyridin-4-yl	H	H	H	(±)- <i>cis</i>
161	B.13	c.b.	2-naphthyl	H	H	H	(±)- <i>cis</i>
162	B.13	c.b.	2-quinoliny	H	H	H	(±)- <i>cis</i>
163	B.13	c.b.	3-isopropoxybenzyl	H	H	H	(±)- <i>cis</i>
164	B.13	c.b.	1-phenylethyl	H	H	H	(±)- <i>cis</i>
165	B.13	c.b.	3-isopropoxyphenyl	H	H	H	(±)- <i>trans</i>
166	B.13	c.b.	3-cyanophenyl	H	H	H	(±)- <i>trans</i>
167	B.13	c.b.		H	H	H	(±)- <i>trans</i>
168	B.13	c.b.	2,4-dichlorophenyl	H	H	H	(±)- <i>trans</i>
169	B.13	c.b.	2-thienyl	H	H	H	(±)- <i>trans</i>

& c.b. = covalent bond

Table 4



Co.No.	Ex.	R <sub>a</sub>	X&	R <sup>2</sup>	L	Physical data
170	B.1	CF <sub>3</sub>	c.b.	3,5-di(trifluoro-methyl)phenyl	H	(±)-(A)
171	B.1	CF <sub>3</sub>	c.b.	3,5-di(trifluoro-methyl)phenyl	H	(±)-(B) fumaric acid (1:4)
172	B.13	H	c.b.	3-trifluoromethyl-phenyl	benzyl	(±)-cis
173	B.13	H	c.b.	3,5-difluoro-phenyl	benzyl	(±)-cis
174	B.13	H	c.b.	2-naphthyl	benzyl	(±)-cis
175	B.13	H	c.b.	3-cyano-phenyl	benzyl	(±)-cis
176	B.13	H	-O-	benzyl	benzyl	(±)-cis
177	B.13	H	c.b.	2-furanyl	benzyl	(±)-cis
178	B.13	H	c.b.	2-thienyl	benzyl	(±)-cis
179	B.13	H	c.b.	phenyl	benzyl	(±)-cis
180	B.13	H	c.b.	3,5-dichloro-phenyl	benzyl	(±)-cis
181	B.17	H	c.b.	3,4-dichlorophenyl	benzyl	(±)-cis
182	B.13	H	c.b.		benzyl	(±)-cis
183	B.13	H	c.b.	phenyl	benzyl	(±)-trans
184	B.13	H	c.b.	2,6-dichloro-pyridin-4-yl	benzyl	(±)-trans
185	B.13	H	c.b.	2-furanyl	benzyl	(±)-trans
186	B.13	H	c.b.	2-thienyl	benzyl	(±)-trans
187	B.13	H	c.b.	3-cyano-phenyl	benzyl	(±)-trans
188	B.13	H	c.b.		benzyl	(±)-trans
189	B.13	H	c.b.		benzyl	(±)-trans
190	B.13	H	c.b.	5-methyl-isoxazol-3-yl	benzyl	(±)-trans
191	B.13	H	c.b.	2-nitrophenyl	benzyl	(±)-cis
192	B.16	H	c.b.	2-aminophenyl	benzyl	(±)-cis fumaric acid (1:2)

& c.b. = covalent bond

[0092] Table 5 lists both the experimental (column heading "Exp") and theoretical (column heading "Theor") elemental analysis values for carbon, hydrogen and nitrogen for the compounds as prepared in the experimental part hereinabove.

Table 5

Co. No.	C		H		N	
	Exp	Theor	Exp	Theor	Exp	Theor
5	75.47	76.05	8.06	8.02	9.91	10.14
6	75.08	76.05	8.18	8.02	10.02	10.14
7	68.40	68.18	6.68	6.68	5.52	5.68

EP 0 862 566 B1

Table 5 (continued)

		C		H		N	
	Co. No.	Exp	Theor	Exp	Theor	Exp	Theor
5	8	67.39	68.18	6.83	6.68	5.62	5.68
	9	71.98	71.77	7.59	7.40	7.11	7.17
	10	76.99	77.23	8.12	7.90	8.16	8.44
	12	81.31	81.83	7.79	7.77	7.36	7.53
10	16	64.88	65.19	5.53	5.64	7.05	7.13
	17	58.45	58.46	4.85	5.03	5.01	5.11
	19	61.61	61.91	5.03	5.35	10.32	10.62
	20	60.51	60.71	4.92	4.92	6.89	7.08
15	21	60.74	60.71	4.86	4.92	7.02	7.08
	22	63.09	62.87	5.56	5.72	10.08	10.18
	23	62.68	62.98	5.21	5.28	11.43	11.60
	24	62.44	62.98	5.18	5.28	11.33	11.60
20	25	60.33	61.53	5.42	5.74	11.62	11.96
	26	63.42	63.63	5.96	5.80	8.52	8.48
	27	63.41	63.63	5.82	5.80	8.04	8.48
	30	63.02	62.94	5.19	5.28	7.01	7.10
25	31	62.68	62.94	5.31	5.28	7.08	7.10
	32	66.40	66.47	5.44	5.58	7.66	7.75
	33	66.42	66.47	5.60	5.58	7.55	7.75
	34	62.07	62.98	5.10	5.28	11.46	11.60
30	35	65.11	65.04	5.08	5.03	5.89	5.99
	36	64.94	65.04	4.97	5.03	6.03	5.99
	37	62.58	62.65	5.26	5.42	8.63	8.86
	38	60.87	61.62	5.22	5.48	8.23	8.45
35	40	62.28	62.43	4.78	5.08	11.28	11.38
	41	63.44	63.63	5.69	5.80	8.36	8.48
	42	63.83	63.63	5.77	5.80	8.40	8.48
	43	66.54	66.75	5.52	5.60	5.61	5.70
40	45	56.47	56.50	4.40	4.60	7.89	7.99
	46	64.49	64.71	5.77	5.87	7.92	8.16
	47	63.46	63.63	5.84	5.80	8.39	8.48
	48	65.16	65.53	6.38	6.20	7.71	7.84
45	49	65.43	65.53	6.25	6.20	7.81	7.84
	50	63.71	63.63	5.64	5.80	8.40	8.48
	51	61.79	61.62	5.46	5.48	8.31	8.45
	52	64.50	64.71	5.70	5.87	7.92	8.16
50	53	63.77	63.63	5.90	5.80	8.25	8.48
	54	63.54	64.53	5.15	5.29	5.19	5.37
	55	67.97	68.56	5.65	5.60	6.02	6.31
	56	63.34	63.46	5.39	5.49	6.79	6.94
55	57	63.54	63.46	5.43	5.49	6.97	6.94
	59	63.58	63.63	5.75	5.80	8.37	8.48
	60	64.13	63.97	5.61	5.69	6.60	6.78
	61	63.41	63.97	5.65	5.69	6.60	6.78
60	62	62.65	62.87	5.76	5.72	9.97	10.18
	63	58.77	58.92	5.14	5.19	4.99	5.03
	65	63.78	64.08	6.24	5.98	7.97	8.30
	66	62.89	64.22	5.41	5.39	11.03	10.96

EP 0 862 566 B1

Table 5 (continued)

	Co. No.	C		H		N	
		Exp	Theor	Exp	Theor	Exp	Theor
5	67	63.06	64.22	5.18	5.39	10.62	10.96
	69	60.74	61.28	5.57	5.68	11.15	11.28
	70	62.34	62.53	4.81	5.11	11.62	11.82
	71	54.59	54.69	4.03	4.20	7.04	7.29
10	72	59.45	59.57	5.01	5.28	9.71	9.92
	74	54.84	54.85	4.18	4.47	9.28	9.41
	76	59.66	60.19	6.21	6.06	8.93	9.36
	78	63.95	64.08	5.90	5.98	8.23	8.30
	79	55.18	55.25	4.08	4.38	7.04	7.16
15	82	62.65	63.63	5.80	5.80	8.22	8.48
	83	61.84	63.63	5.91	5.80	8.00	8.48
	84	58.20	58.49	5.21	5.38	12.81	13.20
	85	61.55	62.53	5.15	5.11	11.53	11.82
20	86	63.82	63.97	5.58	5.69	6.73	6.78
	87	67.38	68.37	5.48	6.15	5.48	5.70
	92	61.87	61.95	5.17	5.32	6.32	6.72
	93	57.14	57.47	4.71	4.92	5.02	5.26
	94	66.62	66.84	5.55	5.75	7.23	7.60
25	95	58.94	58.94	5.60	5.58	6.97	7.05
	96	52.25	52.35	4.82	4.97	11.15	11.25
	97	62.54	62.35	5.26	5.37	11.64	11.79
	98	63.86	63.82	5.30	5.51	8.39	8.51
30	99	64.29	64.28	5.40	5.69	8.17	8.33
	100	62.13	62.35	5.16	5.37	11.59	11.79
	101	64.49	64.28	5.68	5.69	8.08	8.33
	102	67.17	66.84	5.82	5.75	7.36	7.60
	103	58.42	58.69	5.39	5.47	6.15	6.08
35	104	64.16	64.28	5.73	5.69	8.31	8.33
	105	63.95	64.08	6.01	5.98	8.25	8.30
	106	62.27	62.65	5.37	5.42	8.84	8.86
	107	63.76	63.82	5.54	5.51	8.45	8.51
40	108	54.46	54.69	4.08	4.20	7.20	7.29
	109	58.46	58.49	5.19	5.38	12.90	13.20
	110	56.57	57.14	4.58	4.80	7.96	8.13
	111	56.80	57.14	4.59	4.80	7.99	8.13
	112	56.42	56.33	4.73	4.88	10.77	10.95
45	114	54.52	54.17	4.61	4.82	5.75	5.74
	115	64.06	64.08	6.07	5.98	8.16	8.30
	116	63.87	64.08	5.89	5.98	8.11	8.30
	117	55.13	55.25	4.18	4.38	7.05	7.16
50	118	57.96	58.41	4.90	5.12	6.06	6.19
	119	60.60	60.60	5.45	5.42	9.12	9.42
	120	59.92	59.72	4.73	4.70	6.20	6.53
	121	63.51	63.82	5.60	5.51	8.43	8.51
	122	63.53	63.82	5.77	5.51	9.05	8.51
55	123	64.81	65.00	5.75	5.75	10.17	10.24
	131	63.23	63.15	5.72	5.61	8.51	8.66
	132	60.00	60.34	4.90	5.21	7.76	8.04

Table 5 (continued)

Co. No.	C		H		N	
	Exp	Theor	Exp	Theor	Exp	Theor
133	59.93	60.34	5.17	5.21	7.94	8.04
134	60.22	60.34	5.22	5.21	7.94	8.04
135	60.87	60.34	5.44	5.21	7.97	8.04
137	59.04	59.34	5.01	5.12	7.43	7.69
138	57.81	57.62	4.76	4.97	7.54	7.68
139	57.28	57.62	4.73	4.97	7.25	7.68
140	57.53	57.91	4.88	4.86	7.18	7.50
141	57.45	57.91	4.79	4.86	7.21	7.50
142	73.32	74.19	8.25	7.96	9.28	9.61
143	61.66	62.21	5.77	5.82	10.21	10.36
144	75.31	75.54	7.76	7.68	10.28	10.68
145	76.49	77.32	7.43	7.37	9.33	9.75
170	53.73	55.03	4.46	4.62	7.14	7.40
192	64.48	65.13	6.20	6.33	7.84	7.99

C. Pharmacological exampleExample C.1 : Antagonism of substance-P induced relaxation of the pig coronary arteries

[0093] Segments of coronary arteries taken from pigs (killed by injection of an overdose of sodium pentobarbital) were inverted and mounted for recording of isometric tension in organ baths (volume 20 ml) with the endothelium at the outside. The preparations were bathed in Krebs - Henseleit solution. The solution was kept at 37 °C and gassed with a mixture of O<sub>2</sub>/CO<sub>2</sub> (95/5). After stabilisation of the preparations, prostaglandin F<sub>2α</sub> (10<sup>-5</sup> M) was administered to induce a contraction. This was repeated until contractile responses became stable. Then prostaglandin F<sub>2α</sub> was again administered and substance P (3x10<sup>-10</sup> M and 10<sup>-9</sup> M cumulatively) was added. Substance P induced endothelium dependent relaxations. After washing away the agonists, a known concentration of a compound of formula (I) was added. After an incubation period of 30 minutes, prostaglandin F<sub>2α</sub> (10<sup>-5</sup> M) and the same concentrations of substance P as described above were again administered in the presence of the compound to be tested. Relaxations caused by substance P were expressed as relaxations under control conditions, and percent inhibition of the response to 10<sup>-9</sup> M substance P was taken as a measure of the antagonistic activity of the compound to be tested. Table 6 lists the IC<sub>50</sub> values (concentration at which 50 % of the response to 10<sup>-9</sup> M substance P was inhibited by the test compound) for the tested compounds.



Table 6

Co. No.	IC <sub>50</sub> (in 10 <sup>-9</sup> M)	Co. No.	IC <sub>50</sub> (in 10 <sup>-9</sup> M)	Co. No.	IC <sub>50</sub> (in 10 <sup>-9</sup> M)
5	4.61	51	1.35	97	1.58
17	1.68	52	1.10	99	1.46
19	0.54	53	0.35	101	0.75
22	0.37	55	2.8	102	1.85
24	0.64	56	3.25	104	0.30
25	0.79	58	0.24	105	1.05
26	2.75	59	1.20	107	0.96
27	0.13	61	15.2	115	2.23
28	13.3	62	0.24	119	2.81
29	0.45	63	8.38	120	0.82
33	0.60	65	6.39	121	2.77
34	0.35	68	5.88	122	1.68
35	17.0	69	1.57	124	0.06
36	2.31	70	0.29	128	0.35
37	9.60	73	5.73	132	2.08
38	0.86	76	14.1	133	0.62
42	0.93	85	0.15	134	0.04
43	5.63	86	2.13	135	0.01
44	8.34	87	1.90	136	0.31
45	0.15	91	0.07	137	0.23
46	0.42	92	0.78	138	0.16
48	0.26	93	4.99	139	0.13
49	0.59	94	0.42		
50	2.43	95	0.10		

Example C.2 : Antagonism of substance P induced plasma extravasation in guinea-pigs

[0094] Plasma extravasation was induced by injection of substance P (2 mg/kg) in the femoral artery of female guinea-pigs. Evans Blue dye (30 mg/kg) was injected simultaneously. The test compound or solvent was administered 1 hour prior to substance P injection. 10 minutes after challenge, the animals were checked for blue colouring (a direct measure for plasma extravasation) of the nose, the forepaws, and the conjunctiva. 30 minutes after challenge, the animals were sacrificed by CO<sub>2</sub> gas inhalation and checked for blue colouring of the trachea and the urinary bladder. Doses which actively inhibit substance P-induced plasma extravasation are defined as those doses at which only 1/3 or less of the total surface area of the nose, forepaws, conjunctiva, trachea or urinary bladder are coloured blue by an intensive extravasation. Table 7 lists the lowest active doses (LAD) in mg/kg for the tested compounds.

Table 7

Co. No.	LAD (in mg/kg)				
	nose	forepaws	conjunctiva	trachea	urinary bladder
8	10.0	10.0	10.0	> 40.0	> 40.0
12	> 40.0	> 40.0	> 40.0	> 40.0	40.0
16	40.0	40.0	40.0	> 40.0	> 40.0
17	10.0	10.0	10.0	> 40.0	10.0
19	2.50	2.50	2.50	10.0	> 40.0
21	10.0	10.0	10.0	> 40.0	10.0
22	2.50	2.50	2.50	10.0	> 40.0
23	10.0	10.0	> 40.0	> 40.0	> 40.0
24	2.50	2.50	10.0	10.0	> 40.0
25	2.50	2.50	10.0	> 40.0	> 40.0
26	10.0	10.0	20.0	40.0	20.0
27	0.63	0.63	0.63	0.63	0.63
33	2.50	10.0	2.50	> 40.0	> 40.0
34	2.50	10.0	2.50	10.0	10.0
38	10.0	10.0	10.0	> 40.0	> 40.0
40	10.0	10.0	10.0	> 40.0	> 40.0
42	1.25	1.25	2.50	5.00	5.00
45	0.63	0.63	0.63	> 40.0	2.50
46	0.63	0.63	0.63	> 40.0	0.63
47	40.0	40.0	40.0	> 40.0	> 40.0
48	2.50	2.50	2.50	2.50	2.50
49	2.50	2.50	2.50	10.0	> 40.0
50	10.0	10.0	10.0	10.0	> 40.0
52	0.63	0.63	2.50	10.0	10.0
53	1.25	1.25	1.25	2.50	2.50
56	10.0	10.0	10.0	40.0	2.50
59	2.50	2.50	2.50	5.00	5.00
68	10.0	10.0	> 40.0	> 40.0	> 40.0
70	10.0	10.0	10.0	10.0	10.0
73	10.0	10.0	10.0	> 40.0	> 40.0
74	10.0	10.0	10.0	> 40.0	> 40.0
82	2.50	2.50	2.50	10.0	> 40.0
83	0.63	0.63	2.50	> 40.0	0.63
85	2.50	2.50	2.50	2.50	10.0
87	10.0	10.0	10.0	10.0	> 40.0
90	10.0	10.0	10.0	10.0	> 40.0
94	2.50	2.50	2.50	10.0	10.0
95	0.31	0.31	0.31	0.63	2.50
96	10.0	10.0	10.0	> 40.0	> 40.0
101	10.0	10.0	10.0	> 40.0	> 40.0
103	2.50	2.50	2.50	10.0	2.50
105	2.50	2.50	2.50	> 40.0	> 40.0
107	2.50	2.50	2.50	2.50	10.0
119	2.50	2.50	2.50	10.0	> 40.0
128	0.63	0.63	0.63	> 40.0	10.0
132	10.0	10.0	10.0	10.0	10.0
133	2.50	2.50	2.50	10.0	10.0

Table 7 (continued)

Co. No.	LAD (in mg/kg)				
	nose	forepaws	conjunctiva	trachea	urinary bladder
134	10.0	10.0	10.0	10.0	10.0
135	0.63	0.63	0.63	2.50	> 40.0
136	2.50	2.50	2.50	2.50	2.50
137	2.50	2.50	2.50	2.50	2.50
138	2.50	2.50	2.50	10.0	> 40.0
139	2.50	2.50	2.50	2.50	2.50

D. Composition examples

[0095] "Active ingredient" (A.I.) as used throughout these examples relates to a compound of formula (I) a pharmaceutically acceptable addition salt, a stereochemically isomeric form thereof or a *N*-oxide form thereof.

Example D.1 : ORAL DROPS

[0096] 500 Grams of the A.I. was dissolved in 0.5 l of 2-hydroxypropanoic acid and 1.5 l of the polyethylene glycol at 60~80°C. After cooling to 30~40°C there were added 35 l of polyethylene glycol and the mixture was stirred well. Then there was added a solution of 1750 grams of sodium saccharin in 2.5 l of purified water and while stirring there were added 2.5 l of cocoa flavor and polyethylene glycol q.s. to a volume of 50 l, providing an oral drop solution comprising 10 mg/ml of A.I. The resulting solution was filled into suitable containers.

Example D.2 : ORAL SOLUTION

[0097] 9 Grams of methyl 4-hydroxybenzoate and 1 gram of propyl 4-hydroxybenzoate were dissolved in 4 l of boiling purified water. In 3 l of this solution were dissolved first 10 grams of 2,3-dihydroxybutanedioic acid and thereafter 20 grams of the A.I. The latter solution was combined with the remaining part of the former solution and 12 l 1,2,3-propanetriol and 3 l of sorbitol 70% solution were added thereto. 40 Grams of sodium saccharin were dissolved in 0.5 l of water and 2 ml of raspberry and 2 ml of gooseberry essence were added. The latter solution was combined with the former, water was added q.s. to a volume of 20 l providing an oral solution comprising 5 mg of the active ingredient per teaspoonful (5 ml). The resulting solution was filled in suitable containers.

Example D.3 : FILM-COATED TABLETSPreparation of tablet core

[0098] A mixture of 100 grams of the A.I., 570 grams lactose and 200 grams starch was mixed well and thereafter humidified with a solution of 5 grams sodium dodecyl sulfate and 10 grams polyvinylpyrrolidone in about 200 ml of water. The wet powder mixture was sieved, dried and sieved again. Then there was added 100 grams microcrystalline cellulose and 15 grams hydrogenated vegetable oil. The whole was mixed well and compressed into tablets, giving 10.000 tablets, each containing 10 mg of the active ingredient.

Coating

[0099] To a solution of 10 grams methyl cellulose in 75 ml of denaturated ethanol there was added a solution of 5 grams of ethyl cellulose in 150 ml of dichloromethane. Then there were added 75 ml of dichloromethane and 2.5 ml 1,2,3-propanetriol. 10 Grams of polyethylene glycol was molten and dissolved in 75 ml of dichloromethane. The latter solution was added to the former and then there were added 2.5 grams of magnesium octadecanoate, 5 grams of polyvinylpyrrolidone and 30 ml of concentrated colour suspension and the whole was homogenated. The tablet cores were coated with the thus obtained mixture in a coating apparatus.


Example D.4 : INJECTABLE SOLUTION

[0100] 1.8 Grams methyl 4-hydroxybenzoate and 0.2 grams propyl 4-hydroxybenzoate were dissolved in about 0.5 l of boiling water for injection. After cooling to about 50°C there were added while stirring 4 grams lactic acid, 0.05

grams propylene glycol and 4 grams of the A.I.. The solution was cooled to room temperature and supplemented with water for injection q.s. ad 1 l, giving a solution comprising 4 mg/ml of A.I.. The solution was sterilized by filtration and filled in sterile containers.

# Claims

## 1. A compound of formula



a *N*-oxide form, a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof, wherein

*n* is 0, 1 or 2;

*m* is 1 or 2, provided that if *m* is 2, then *n* is 1;

*p* is 1 or 2;

*=Q* is =O or =NR<sup>3</sup>;





*X* is a covalent bond or a bivalent radical of formula -O-, -S-, -NR<sup>3</sup>-;

*R*<sup>1</sup> is Ar<sup>1</sup>, Ar<sup>1</sup>C<sub>1-6</sub>alkyl or di(Ar<sup>1</sup>)C<sub>1-6</sub>alkyl, wherein each C<sub>1-6</sub>alkyl group is optionally substituted with hydroxy, C<sub>1-4</sub>alkyloxy, oxo or a ketalized oxo substituent of formula -O-CH<sub>2</sub>-CH<sub>2</sub>-O- or -O-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O-;


*R*<sup>2</sup> is Ar<sup>2</sup>, Ar<sup>2</sup>C<sub>1-6</sub>alkyl, Het<sup>1</sup> or Het<sup>1</sup>C<sub>1-6</sub>alkyl;

*R*<sup>3</sup> is hydrogen or C<sub>1-6</sub>alkyl;

*L* is hydrogen; Ar<sup>3</sup>; C<sub>1-6</sub>alkyl; C<sub>1-6</sub>alkyl substituted with 1 or 2 substituents selected from hydroxy, C<sub>1-6</sub>alkyloxy, Ar<sup>3</sup>, Ar<sup>3</sup>C<sub>1-6</sub>alkyloxy and Het<sup>2</sup>; C<sub>3-6</sub>alkenyl; Ar<sup>3</sup>C<sub>3-6</sub>alkenyl; di(Ar<sup>3</sup>)C<sub>3-6</sub>alkenyl or a radical of formula



or



wherein

- each  $q$  independently is 2, 3 or 4;  
 each  $r$  is 0, 1, 2, 3 or 4;  
 each  $\text{Y}^1$  independently is a covalent bond,  $-\text{O}-$  or  $\text{NR}^3$ ;  
 $\text{Y}^2$  is a covalent bond,  $\text{C}_{1-4}$ alkanediyl or  $-\text{C}_{1-4}\text{alkylNR}^3$ ;  
 each  $-\text{A}=\text{B}-$  independently is a bivalent radical of formula  $-\text{CH}=\text{CH}-$ ,  $-\text{N}=\text{CH}-$  or  $-\text{CH}=\text{N}-$ ;  
 each  $\text{R}^4$  independently is hydrogen,  $\text{C}_{1-6}$ alkyl,  $\text{Ar}^2$  or  $\text{Ar}^2\text{C}_{1-6}$ alkyl;  
 $\text{R}^5$  is hydrogen,  $\text{C}_{1-6}$ alkyl or  $\text{Ar}^3$ ;  
 $\text{R}^6$  is  $\text{C}_{1-6}$ alkyl,  $\text{Ar}^3$ ,  $\text{Ar}^3\text{C}_{1-6}$ alkyl,  $\text{di}(\text{Ar}^3)\text{C}_{1-6}$ alkyl,  $\text{Ar}^3\text{C}_{3-7}$ cycloalkyl, or indolyl;  
 $\text{R}^7$  is  $\text{Ar}^3$ ,  $\text{Ar}^3\text{C}_{1-6}$ alkyl;  $\text{di}(\text{Ar}^3)\text{C}_{1-6}$ alkyl;  $\text{C}_{1-6}$ alkyl;  $\text{C}_{3-7}$ cycloalkyl;  $\text{C}_{3-7}$ cycloalkyl substituted with  $\text{Ar}^3$ ; oxazolyl; oxazolyl substituted with halo or  $\text{C}_{1-6}$ alkyl; thiazolyl; thiazolyl substituted with halo or  $\text{C}_{1-6}$ alkyl; imidazolyl; imidazolyl substituted with  $\text{Ar}^3$ ,  $\text{C}_{1-6}$ alkyl,  $\text{Ar}^3\text{C}_{1-6}$ alkyl or halo; indolyl; indolyl substituted with  $\text{C}_{1-4}$ alkyl; 2,3,4-trihydroquinolyl; pyrrolidinyl or furanyl;  
 each  $\text{R}^8$  independently is hydrogen,  $\text{C}_{1-6}$ alkyl,  $\text{C}_{3-7}$ cycloalkyl or a radical of formula



or



wherein

- $\text{Alk}$  is  $\text{C}_{1-6}$ alkanediyl;  
 $\text{Z}$  is a bivalent radical of formula  $-\text{O}-$ ,  $-\text{S}-$  or  $-\text{NR}^3$ ;  
 $\text{R}^{11}$  is phenyl; phenyl substituted with 1 or 2 substituents selected from halo,  $\text{C}_{1-6}$ alkyl or  $\text{C}_{1-6}$ alkyloxy; furanyl; furanyl substituted with 1 or 2 substituents selected from  $\text{C}_{1-6}$ alkyl or hydroxy $\text{C}_{1-6}$ alkyl; thienyl; thienyl substituted with 1 or 2 substituents selected from halo or  $\text{C}_{1-6}$ alkyl; oxazolyl; oxazolyl substituted with 1 or 2  $\text{C}_{1-6}$ alkyl substituents; thiazolyl; thiazolyl substituted with 1 or 2  $\text{C}_{1-6}$ alkyl substituents; pyridinyl or pyridinyl substituted with 1 or 2  $\text{C}_{1-6}$ alkyl substituents;  
 $\text{R}^{12}$  is  $\text{C}_{1-6}$ alkyl or  $\text{C}_{1-6}$ alkyl substituted with hydroxy, carboxyl or  $\text{C}_{1-6}$ alkyloxycarbonyl;  
 $\text{Ar}^1$  is phenyl; phenyl substituted with 1, 2 or 3 substituents each independently selected from halo,  $\text{C}_{1-4}$ alkyl, halo $\text{C}_{1-4}$ alkyl, cyano, aminocarbonyl,  $\text{C}_{1-4}$ alkyloxy or halo $\text{C}_{1-4}$ alkyloxy;  
 $\text{Ar}^2$  is naphthalenyl; phenyl; phenyl substituted with 1, 2 or 3 substituents each independently selected from hydroxy, halo, cyano, nitro, amino, mono- or di( $\text{C}_{1-4}$ alkyl)amino,  $\text{C}_{1-4}$ alkyl, halo $\text{C}_{1-4}$ alkyl,  $\text{C}_{1-4}$ alkyloxy, halo $\text{C}_{1-4}$ alkyloxy, carboxyl,  $\text{C}_{1-4}$ alkyloxycarbonyl, aminocarbonyl and mono- or di( $\text{C}_{1-4}$ alkyl)aminocarbonyl;  
 $\text{Ar}^3$  is phenyl or phenyl substituted with 1, 2 or 3 substituents selected from halo, hydroxy, amino, nitro, aminocarbonyl,  $\text{C}_{1-6}$ alkyl, halo $\text{C}_{1-6}$ alkyl or  $\text{C}_{1-6}$ alkyloxy;  
 $\text{Het}^1$  is a monocyclic heterocycle selected from pyrrolyl, pyrazolyl, imidazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl; or a bicyclic heterocycle selected from quinolyl, quinoxalyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl and benzothienyl; each monocyclic and bicyclic heterocycle may optionally be substituted on a carbon atom by 1 or 2 substituents selected from halo,  $\text{C}_{1-4}$ alkyl or mono-, di- or tri(halo)methyl; and

Het<sup>2</sup> is a heterocycle selected from 1,4-dihydro-5-oxo-tetrazol-1-yl, imidazo[1,2-a]pyridinyl, oxazolyl or imidazolyl; each of said heterocycles may be substituted with 1 or where possible 2 substituents selected from C<sub>1-4</sub>alkyl and Ar<sup>3</sup>.

2. A compound according to claim 1 wherein L is hydrogen; C<sub>1-6</sub>alkyl; C<sub>1-6</sub>alkyl substituted with hydroxy; C<sub>3-6</sub>alkenyl; Ar<sup>3</sup>; Ar<sup>3</sup>C<sub>1-6</sub>alkyl; di(Ar<sup>3</sup>)C<sub>1-6</sub>alkyl; Ar<sup>3</sup>C<sub>3-6</sub>alkenyl; di(Ar<sup>3</sup>)C<sub>1-6</sub>alkenyl; or a radical of formula (a-1), (a-2), (a-4) or (a-5) wherein

R<sup>7</sup> is Ar<sup>3</sup>; Ar<sup>3</sup>C<sub>1-6</sub>alkyl; di(Ar<sup>3</sup>)C<sub>1-6</sub>alkyl; C<sub>1-6</sub>alkyl; C<sub>3-7</sub>cycloalkyl; C<sub>3-7</sub>cycloalkyl substituted with Ar<sup>3</sup>; oxazolyl; oxazolyl substituted with halo or C<sub>1-6</sub>alkyl; thiazolyl; thiazolyl substituted with halo or C<sub>1-6</sub>alkyl; imidazolyl; imidazolyl substituted with Ar<sup>3</sup>, C<sub>1-6</sub>alkyl, Ar<sup>3</sup>C<sub>1-6</sub>alkyl or halo; pyrrolidinyl or furanyl;

Ar<sup>3</sup> is is phenyl or phenyl substituted with 1, 2 or 3 substituents selected from halo, hydroxy, amino, aminocarbonyl, C<sub>1-6</sub>alkyl, haloC<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxy;

Het<sup>1</sup> is a monocyclic heterocycle selected from pyrrolyl, pyrazolyl, imidazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl; or a bicyclic heterocycle selected from quinolinyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl and benzothienyl; each monocyclic and bicyclic heterocycle may optionally be substituted on a carbon atom by 1 or 2 substituents selected from halo, C<sub>1-4</sub>alkyl or mono-, di- or tri(halo)methyl.

3. A compound as claimed in claim 1 or 2 wherein R<sup>1</sup> is Ar<sup>1</sup>C<sub>1-6</sub>alkyl, R<sup>2</sup> is phenyl substituted with 2 substituents selected from methyl and trifluoromethyl, X is a covalent bond and =Q is =O.

4. A compound as claimed in any one of claims 1 to 3 wherein n and m are 1 and p is 1 or 2.

5. A compound as claimed in any one of claims 1 to 4 wherein R<sup>1</sup> is phenylmethyl; R<sup>2</sup> is phenyl substituted with 2 substituents selected from methyl or trifluoromethyl; n, m and p are 1; X is a covalent bond; and =Q is =O.

6. A compound as claimed in any one of claims 1 to 4 wherein L is a radical of formula (a-2) wherein R<sup>4</sup> is hydrogen or phenyl; r is 0 or 1; Y<sup>1</sup> is a covalent bond, -O- or -NH-; R<sup>7</sup> is pyrrolidinyl; furanyl; 1-phenylcyclohexanyl; diphenylmethyl; or phenyl substituted with 1, 2 or 3 substituents each independently selected from methyl, methoxy or chloro.

7. A compound as claimed in claim 5 or 6 wherein the compound has the *trans* configuration.

8. A compound as claimed in claim 5 or 6 wherein the compound has the *cis* configuration.

9. A compound as claimed in claim 1, wherein L is hydrogen.

10. A compound as claimed in claim 1, wherein the compound is

4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]-N-(2,6-dimethylphenyl)-1-piperazine acetamide;

4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]-N-(1-phenylcyclohexyl)-1-piperazine acetamide;

1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-[4-[α-(1-pyrrolidinylcarbonyl)benzyl]-1-piperazinyl]piperidine;

1-[3,5-bis(trifluoromethyl)benzoyl]-4-[4-[1-[(2-methyl-5-oxazolyl)methyl]-1H-benzimidazol-2-yl]-1-piperazinyl]-2-(phenylmethyl)piperidine;

4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-[(4-trifluoromethylphenyl)methyl]-4-piperidinyl]-N-(2,6-dimethylphenyl)-1-piperazine acetamide;

4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-N-(2,6-dimethylphenyl)-1-piperazine acetamide.

11. A compound as claimed in claim 10, wherein the compound is


(+)-(B)-*trans*-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]-N-(2,6-dimethylphenyl)-1-piperazine acetamide;

(-)-(B)-*cis*-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]-N-(2,6 dimethylphenyl)-1-pip-

erazine acetamide, or  
 (+)-(B)-*trans*-4-[1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinyl]-*N*-(2,6-dimethylphenyl)-  
 1-piperazine acetamide, (L)-malic acid (1:1).


- 5 12. A composition comprising a pharmaceutically acceptable carrier, and as active ingredient a therapeutically effective amount of a compound as claimed in any one of claims 1 to 11.
13. A process of preparing a composition as claimed in claim 12, characterized in that a pharmaceutically acceptable carrier is intimately mixed with a therapeutically effective amount of a compound as claimed in any one claims 1 to 11.
- 10 14. A compound as claimed in any one of claims 1 to 11 for use as a medicine.
- 15 15. A process of preparing a compound as claimed in claim 1, characterized by

a) reductively *N*-alkylating an intermediate of formula (II) wherein L and p are defined as in claim 1, with an intermediate of formula (I)




wherein R<sup>1</sup>, R<sup>2</sup>, X, Q, n and m are defined as in claim 1, in a reaction-inert solvent, in the presence of an appropriate reducing agent and optionally in the presence of a suitable complex-forming agent;

b) reacting an intermediate of formula (IV) wherein R<sup>2</sup>, X and Q are defined as in claim 1 and W<sup>1</sup> is an appropriate leaving group with an intermediate of formula (V)




wherein R<sup>1</sup>, L, n, m and p are defined as in claim 1, in a reaction-inert solvent and in the presence of a suitable base;

c) reductively *N*-alkylating a piperazine derivative of formula (VII) wherein p is defined as in claim 1 and P<sup>1</sup> is a protective group, with an intermediate of formula (II)




wherein  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{X}$ ,  $\text{Q}$ ,  $n$  and  $m$  are defined as in claim 1, in a reaction-inert solvent, in the presence of an appropriate reducing agent and optionally in the presence of a suitable complex-forming agent, thus forming a compound of formula (I-c);

d) reductively *N*-alkylating a piperazine derivative of formula (VII) wherein  $p$  is defined as in claim 1 and  $\text{P}^1$  is a protective group, with an intermediate of formula (VIII),




wherein  $\text{R}^1$ ,  $n$  and  $m$  are defined as in claim 1, in a reaction-inert solvent and in the presence of a suitable base, thus forming an intermediate of formula (XI); which may subsequently be reacted with an intermediate of formula (IV)




wherein  $\text{W}^1$  is an appropriate leaving group, and  $\text{X}$ ,  $\text{Q}$  and  $\text{R}^2$  are defined as in claim 1, to form a compound of formula (I-c);

e) deprotecting a compound of formula (I-c) using art-known deprotection techniques, thus forming a compound of formula (I-b);





f) reacting a compound of formula (I-b) with an intermediate of formula (VI)




wherein L' is the same as L as defined in claim 1 but other than hydrogen and W<sup>2</sup> is an appropriate leaving group in a reaction-inert solvent and in the presence of a suitable base; thus forming a compound of formula (I-a);

and, if desired, converting compounds of formula (I) into each other following art-known transformations, and further, if desired, converting the compounds of formula (I), into a therapeutically active non-toxic acid addition salt by treatment with an acid, or into a therapeutically active non-toxic base addition salt by treatment with a base, or conversely, converting the acid addition salt form into the free base by treatment with alkali, or converting the base addition salt into the free acid by treatment with acid; and, if desired, preparing stereochemically isomeric forms or N-oxide forms thereof.

## Patentansprüche





### 1. Verbindung der Formel




bzw. ihre N-Oxidform, ihr pharmazeutisch unbedenkliches Additionssalz oder ihre stereochemisch isomere Form, worin

- n 0, 1 oder 2 darstellt,  
 m 1 oder 2 darstellt, unter der Voraussetzung, daß, wenn 2 ist, n 1 ist,  
 p 1 oder 2 darstellt,  
 =Q =O oder =NR<sup>3</sup> darstellt,  
 X eine kovalente Bindung oder einen zweiwertigen Rest der Formel -O-, -S-, -NR<sup>3</sup>- darstellt,  
 R<sup>1</sup> Ar<sup>1</sup>, Ar<sup>1</sup>C<sub>1-6</sub>Alkyl oder Di(Ar<sup>1</sup>)C<sub>1-6</sub>Alkyl darstellt, wobei jede C<sub>1-6</sub>Alkylgruppe gegebenenfalls durch Hydroxy, C<sub>1-4</sub>Alkyloxy, Oxo oder einen ketalisierten Oxosubstituenten der Formel -O-CH<sub>2</sub>-CH<sub>2</sub>-O- oder -O-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O- substituiert ist,  
 R<sup>2</sup> Ar<sup>2</sup>, Ar<sup>2</sup>C<sub>1-6</sub>Alkyl, Het<sup>1</sup> oder Het<sup>1</sup>C<sub>1-6</sub>Alkyl darstellt;  
 R<sup>3</sup> Wasserstoff oder C<sub>1-6</sub>Alkyl darstellt,  
 L Wasserstoff, Ar<sup>3</sup>, C<sub>1-6</sub>Alkyl, C<sub>1-6</sub>Alkyl, das durch 1 oder 2 Substituenten aus der Gruppe Hydroxy, C<sub>1-6</sub>Alkyloxy, Ar<sup>3</sup>, Ar<sup>3</sup>C<sub>1-6</sub>Alkyloxy und Het<sup>2</sup> substituiert ist, C<sub>3-6</sub>Alkenyl, Ar<sup>3</sup>C<sub>3-6</sub>Alkenyl, Di(Ar<sup>3</sup>)

C<sub>3-6</sub>Alkenyl oder einen Rest der Formel



oder



darstellt,  
wobei

q jeweils unabhängig voneinander 2, 3 oder 4 darstellt,

r jeweils 0, 1, 2, 3 oder 4 darstellt,

Y<sup>1</sup> jeweils unabhängig voneinander eine kovalente Bindung, -O- oder NR<sup>3</sup> darstellt,

Y<sup>2</sup> eine kovalente Bindung, C<sub>1-4</sub>Alkandyl oder C<sub>1-4</sub>AlkylNR<sup>3</sup>- darstellt,

-A=B- jeweils unabhängig voneinander einen zweiwertigen Rest der Formel -CH=CH-, -N=CH- oder -CH=N- darstellt,

R<sup>4</sup> jeweils unabhängig voneinander Wasserstoff, C<sub>1-6</sub>Alkyl, Ar<sup>2</sup> oder Ar<sup>2</sup>C<sub>1-6</sub>Alkyl darstellt,

R<sup>5</sup> Wasserstoff, C<sub>1-6</sub>Alkyl oder Ar<sup>3</sup> darstellt,

R<sup>6</sup> C<sub>1-6</sub>Alkyl, Ar<sup>3</sup>, Ar<sup>3</sup>C<sub>1-6</sub>Alkyl, Di(Ar<sup>3</sup>)C<sub>1-6</sub>Alkyl, Ar<sup>3</sup>C<sub>3-7</sub>Cycloalkyl oder Indolyl darstellt,

R<sup>7</sup> Ar<sup>3</sup>, Ar<sup>3</sup>C<sub>1-6</sub>Alkyl, Di(Ar<sup>3</sup>)C<sub>1-6</sub>Alkyl, C<sub>1-6</sub>Alkyl, C<sub>3-7</sub>Cycloalkyl, C<sub>3-7</sub>Cycloalkyl, das durch Ar<sup>3</sup> substituiert ist, Oxazolyl, Oxazolyl, das durch Halogen oder durch C<sub>1-6</sub>Alkyl substituiert ist, Thiazolyl, Thiazolyl, das durch Halogen oder C<sub>1-6</sub>Alkyl substituiert ist, Imidazolyl, Imidazolyl, das durch Ar<sup>3</sup>, C<sub>1-6</sub>Alkyl, Ar<sup>3</sup>-C<sub>1-6</sub>Alkyl oder Halogen substituiert ist, Indolyl, Indolyl, das durch C<sub>1-4</sub>Alkyl substituiert ist, 2,3,4-Trihydrochinolyl, Pyrrolidinyl oder Furanyl darstellt,

R<sup>8</sup> jeweils unabhängig voneinander Wasserstoff, C<sub>1-6</sub>Alkyl, C<sub>3-7</sub>Cycloalkyl oder einen Rest der Formel



oder

- darstellt,  
 in der Alk C<sub>1-6</sub>Alkandiy l darstellt,  
 Z einen zweiwertigen Rest der Formel -O-, -S- oder -NR<sup>3</sup>- darstellt,  
 R<sup>11</sup> Phenyl, Phenyl, das durch 1 oder 2 Substituenten aus der Gruppe Halogen, C<sub>1-6</sub>Alkyl oder C<sub>1-6</sub>Alkyloxy substituiert ist, Furanyl, Furanyl, das durch 1 oder 2 Substituenten aus der Gruppe C<sub>1-6</sub>Alkyl oder Hydroxy-C<sub>1-6</sub>Alkyl substituiert ist, Thienyl, Thienyl, das durch 1 oder 2 Substituenten aus der Gruppe Halogen oder C<sub>1-6</sub>Alkyl substituiert ist, Oxazolyl, Oxazolyl, das durch 1 oder 2 C<sub>1-6</sub>Alkyl-Substituenten substituiert ist, Thiazolyl, Thiazolyl, das durch 1 oder 2 C<sub>1-6</sub>Alkyl-Substituenten substituiert ist, Pyridinyl, oder Pyridinyl, das durch 1 oder 2 C<sub>1-6</sub>Alkyl-Substituenten substituiert ist, darstellt,  
 R<sup>12</sup> C<sub>1-6</sub>Alkyl oder C<sub>1-6</sub>Alkyl, das durch Hydroxy, Carboxyl oder C<sub>1-6</sub>Alkyloxy-carbonyl substituiert ist, darstellt,  
 Ar<sup>1</sup> Phenyl, Phenyl, das durch 1, 2 oder 3 unabhängige Substituenten aus der Gruppe Halogen, C<sub>1-4</sub>Alkyl, Halogen-C<sub>1-4</sub>Alkyl, Cyan, Aminocarbonyl, C<sub>1-4</sub>Alkyloxy oder Halogen-C<sub>1-4</sub>Alkyloxy substituiert ist, darstellt,  
 Ar<sup>2</sup> Naphthalenyl, Phenyl, Phenyl, das durch 1, 2 oder 3 unabhängige Substituenten aus der Gruppe Hydroxy, Halogen, Cyan, Nitro, Amino, Mono- oder Di(C<sub>1-4</sub>Alkyl) amino, C<sub>1-4</sub>Alkyl, Halogen-C<sub>1-4</sub>Alkyl, C<sub>1-4</sub>Alkyloxy, Halogen-C<sub>1-4</sub>Alkyloxy, Carboxyl, C<sub>1-4</sub>Alkyloxy-carbonyl, Aminocarbonyl und Mono- oder Di(C<sub>1-4</sub>Alkyl)aminocarbonyl darstellt,  
 Ar<sup>3</sup> Phenyl oder Phenyl, das durch 1, 2 oder 3 Substituenten aus der Gruppe Halogen, Hydroxy, Amino, Nitro, Aminocarbonyl, C<sub>1-6</sub>Alkyl, Halogen-C<sub>1-6</sub>alkyl oder C<sub>1-6</sub>Alkyloxy substituiert ist, darstellt,  
 Het<sup>1</sup> einen monozyklischen Heterozyklus aus der Gruppe Pyrrolyl, Pyrazolyl, Imidazolyl, Furanyl, Thienyl, Oxazolyl, Isoxazolyl, Thiazolyl, Isothiazolyl, Pyridinyl, Pyrimidinyl, Pyrazinyl und Pyridazinyl, oder einen bityklischen Heterozyklus aus der Gruppe Chinolinyl, Chinoxalinyl, Indolyl, Benzimidazolyl, Benzoxazolyl, Benzisoxazolyl, Benzothiazolyl, Benzisothiazolyl, Benzofuranyl und Benzothienyl darstellt, wobei jeder monozyklische und bityklische Heterozyklus gegebenenfalls an einem Kohlenstoffatom durch 1 oder 2 Substituenten aus der Gruppe Halogen, C<sub>1-4</sub>Alkyl oder Mono-, Di- oder Tri(halogen)methyl substituiert sein kann, und  
 Het<sup>2</sup> einen Heterozyklus aus der Gruppe 1,4-Dihydro-5-oxo-tetrazol-1-yl, Imidazo[1,2-a]pyridinyl, Oxazolyl oder Imidazolyl darstellt, wobei jeder dieser Heterozyklen durch 1 oder gegebenenfalls 2 Substituenten aus der Gruppe C<sub>1-4</sub>Alkyl und Ar<sup>3</sup> substituiert sein kann.
2. Verbindung nach Anspruch 1, in der L Wasserstoff, C<sub>1-6</sub>Alkyl, C<sub>1-6</sub>Alkyl, das durch Hydroxy substituiert ist, C<sub>3-6</sub>Alkenyl, Ar<sup>3</sup>, Ar<sup>3</sup>C<sub>1-6</sub>Alkyl, Di (Ar<sup>3</sup>) C<sub>1-6</sub>Alkyl, Ar<sup>3</sup>C<sub>3-6</sub>Alkenyl, Di (Ar<sup>3</sup>)C<sub>1-6</sub>Alkenyl oder einen Rest der Formel (a-1), (a-2), (a-4) oder (a-5) darstellt, in dem
- R<sup>7</sup> Ar<sup>3</sup>, Ar<sup>3</sup>C<sub>1-6</sub>Alkyl, Di(Ar<sup>3</sup>)C<sub>1-6</sub>Alkyl, C<sub>1-6</sub>Alkyl, C<sub>3-7</sub>Cycloalkyl, C<sub>3-7</sub>Cycloalkyl, das durch Ar<sup>3</sup> substituiert ist, Oxazolyl, Oxazolyl, das durch Halogen oder C<sub>1-6</sub>Alkyl substituiert ist, Thiazolyl, Thiazolyl, das durch Halogen oder C<sub>1-6</sub>Alkyl substituiert ist, Imidazolyl, Imidazolyl, das durch Ar<sup>3</sup>, C<sub>1-6</sub>Alkyl, Ar<sup>3</sup>-C<sub>1-6</sub>Alkyl oder Halogen substituiert ist, Pyrrolidinyl oder Furanyl darstellt,  
 Ar<sup>3</sup> Phenyl oder Phenyl, das durch 1, 2 oder 3 Substituenten aus der Gruppe Halogen, Hydroxy, Amino, Aminocarbonyl, C<sub>1-6</sub>Alkyl, Halogen-C<sub>1-6</sub>Alkyl oder C<sub>1-6</sub>Alkyloxy substituiert ist, darstellt,  
 Het<sup>1</sup> einen monozyklischen, Heterozyklus aus der Gruppe Pyrrolyl, Pyrazolyl, Imidazolyl, Furanyl, Thienyl, Oxazolyl, Isoxazolyl, Thiazolyl, Isothiazolyl, Pyridinyl, Pyrimidinyl, Pyrazinyl und Pyridazinyl, oder einen bityklischen Heterozyklus aus der Gruppe Chinolinyl, Benzimidazolyl, Benzoxazolyl, Benzisoxazolyl, Benzothiazolyl, Benzisothiazolyl, Benzofuranyl und Benzothienyl darstellt, wobei jeder monozyklische und bityklische Heterozyklus gegebenenfalls an einem Kohlenstoffatom durch 1 oder 2 Substituenten aus der Gruppe Halogen, C<sub>1-4</sub>Alkyl oder Mono-, Di- oder Tri(halogen)methyl substituiert sein kann.
3. Verbindung nach Anspruch 1 oder 2, worin R<sup>1</sup> Ar<sup>1</sup>C<sub>1-6</sub>Alkyl darstellt, R<sup>2</sup> Phenyl, das durch 2 Substituenten aus der Gruppe Methyl und Trifluormethyl substituiert ist, darstellt, X eine kovalente Bindung darstellt und =Q =O darstellt.
4. Verbindung nach einem der Ansprüche 1 bis 3, worin n und m 1 darstellen und p 1 oder 2 darstellt.
5. Verbindung nach einem der Ansprüche 1 bis 4, worin R<sup>1</sup> Phenylmethyl darstellt, R<sup>2</sup> Phenyl, das durch 2 Substi-

tuenten aus der Gruppe Methyl oder Trifluormethyl substituiert ist, darstellt, n, m und p 1 darstellen, X eine kovalente Bindung darstellt und =Q =O darstellt.

6. Verbindung nach einem der Ansprüche 1 bis 4, worin L einen Rest der Formel (a-2) darstellt, in dem R<sup>4</sup> Wasserstoff oder Phenyl darstellt, r 0 oder 1 darstellt, Y<sup>1</sup> eine kovalente Bindung, -O- oder -NH- darstellt, R<sup>7</sup> Pyrrolidinyl, Furanyl, 1-Phenylcyclohexanyl, Diphenylmethyl oder auch Phenyl, das durch 1, 2 oder 3 unabhängige Substituenten aus der Gruppe Methyl, Methoxy oder Chlor substituiert ist, darstellt.

7. Verbindung nach Anspruch 5 oder 6, worin die Verbindung in der trans-Konfiguration vorliegt.

8. Verbindung nach Anspruch 5 oder 6, worin die Verbindung in der cis-Konfiguration vorliegt.

9. Verbindung nach Anspruch 1, worin L Wasserstoff darstellt.

10. Verbindung nach Anspruch 1, wobei es sich bei der Verbindung um

4-[1-[3,5-Bis(trifluormethyl)benzoyl]-2-(phenylmethyl)-4-piperidiny]-N-(2,6-dimethylphenyl)-1-piperazinacetamid;

4-[1-[3,5-Bis(trifluormethyl)benzoyl]-2-(phenylmethyl)-4-piperidiny]-N-(1-phenylcyclohexyl)-1-piperazinacetamid;

1-[3,5-Bis(trifluormethyl)benzoyl]-2-(phenylmethyl)-4-[4-[α-(1-pyrrolidinylcarbonyl)benzyl]-1-piperazinyl]piperidin;

1-[3,5-Bis(trifluormethyl)benzoyl]-4-[4-[1-[(2-methyl-5-oxazolyl)methyl]-1H-benzimidazol-2-yl]-1-piperazinyl]-2-(phenylmethyl)piperidin;

4-[1-[3,5-Bis(trifluormethyl)benzoyl]-2-[(4-trifluormethylphenyl)methyl]-4-piperidiny]-N-(2,6-dimethylphenyl)-1-piperazinacetamid;

4-[1-[3,5-Bis(trifluormethyl)benzoyl]-2-[(3,4-dichlorphenyl)methyl]-4-piperidiny]-N-(2,6-dimethylphenyl)-1-piperazinacetamid handelt.

11. Verbindung nach Anspruch 10, wobei es sich bei der Verbindung um

(+)-(B)-trans-4-[1-[3,5-bis(trifluormethyl)benzoyl]-2-(phenylmethyl)-4-piperidiny]-N-(2,6-dimethylphenyl)-1-piperazinacetamid;

(-)-(B)-cis-4-[1-[3,5-bis(trifluormethyl)benzoyl]-2-(phenylmethyl)-4-piperidiny]-N-(2,6-dimethylphenyl)-1-piperazinacetamid; oder

(+)-(B)-trans-4-[1-[3,5-bis(trifluormethyl)benzoyl]-2-(phenylmethyl)-4-piperidiny]-N-(2,6-dimethylphenyl)-1-piperazinacetamid, (L)-Äpfelsäure (1:1) handelt.


12. Zusammensetzung mit einem pharmazeutisch unbedenklichen Träger und einer therapeutisch wirksamen Menge einer Verbindung nach einem der Ansprüche 1 bis 11 als Wirkstoff.

13. Verfahren zur Herstellung einer Zusammensetzung nach Anspruch 12, dadurch gekennzeichnet, daß man den pharmazeutisch unbedenklichen Träger innig mit einer therapeutisch wirksamen Menge einer Verbindung nach einem der Ansprüche 1 bis 11 mischt.

14. Verbindung nach einem der Ansprüche 1 bis 11 als Arzneimittel.


15. Verfahren zur Herstellung einer Verbindung nach Anspruch 1, dadurch gekennzeichnet, daß man

a) ein Zwischenprodukt der Formel (III) in der L und p wie in Anspruch 1 definiert sind, mit einem Zwischenprodukt der Formel (II)




in der  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{X}$ ,  $\text{Q}$ ,  $n$  und  $m$  wie in Anspruch 1 definiert sind, in einem gegenüber der Reaktion inerten Lösungsmittel in Gegenwart eines geeigneten Reduktionsmittels und gewünschtenfalls in Gegenwart eines geeigneten Komplexbildners reaktiv  $\text{N}$ -alkyliert,

b) ein Zwischenprodukt der Formel (IV), in der  $\text{R}^2$ ,  $\text{X}$  und  $\text{Q}$  wie in Anspruch 1 definiert sind und  $\text{W}^1$  eine geeignete Abgangsgruppe darstellt, mit einem Zwischenprodukt der Formel (V)




in der  $\text{R}^1$ ,  $\text{L}$ ,  $n$ ,  $m$  und  $p$  wie in Anspruch 1 definiert sind, in einem gegenüber der Reaktion inerten Lösungsmittel und in Gegenwart einer geeigneten Base umsetzt,

c) ein Piperazinderivat der Formel (VII), in der  $p$  wie in Anspruch 1 definiert ist und  $\text{P}^1$  eine Schutzgruppe darstellt, mit einem Zwischenprodukt der Formel (II)




in der  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{X}$ ,  $\text{Q}$ ,  $n$  und  $m$  wie in Anspruch 1 definiert sind, in einem gegenüber der Reaktion inerten Lösungsmittel in Gegenwart eines geeigneten Reduktionsmittels und gegebenenfalls in Gegenwart eines geeigneten Komplexbildners reaktiv  $\text{N}$ -alkyliert, wodurch man eine Verbindung der Formel (I-c) erhält,

d) ein Piperazinderivat der Formel (VII), in der  $p$  wie in Anspruch 1 definiert ist und  $\text{P}^1$  eine Schutzgruppe darstellt, mit einem Zwischenprodukt der Formel (VIII),




in der  $\text{R}^1$ ,  $n$  und  $m$  wie in Anspruch 1 definiert sind, in einem gegenüber der Reaktion inerten Lösungsmittel und in Gegenwart einer geeigneten Base reaktiv  $N$ -alkyliert, wodurch man ein Zwischenprodukt der Formel (XI) erhält, das anschließend mit einem Zwischenprodukt der Formel (IV),




in der  $\text{W}^1$  eine geeignete Abgangsgruppe darstellt und  $\text{X}$ ,  $\text{Q}$  und  $\text{R}^2$  wie in Anspruch 1 definiert sind, zu einer Verbindung der Formel (I-c) umgesetzt werden kann,

e) eine Verbindung der Formel (I-c) nach fachbekannten Entschützungsverfahren entschützt, wodurch man eine Verbindung der Formel (I-b) erhält,



f) eine Verbindung der Formel (I-b) mit einem Zwischenprodukt der Formel (VI)




in der  $\text{L}'$  die gleiche Bedeutung wie für  $\text{L}$  in Anspruch 1 definiert hat, jedoch nicht Wasserstoff darstellt, und  $\text{W}^2$  eine geeignete Abgangsgruppe darstellt, in einem gegenüber der Reaktion inerten Lösungsmittel in Gegenwart einer geeigneten Base umgesetzt, wodurch man eine Verbindung der Formel (I-a) erhält,

und gewünschtenfalls Verbindungen der Formel (I) nach fachbekannten Umwandlungsverfahren ineinander umwandelt und weiterhin gewünschtenfalls die Verbindungen der Formel (I) durch Behandeln mit einer Säure in ein therapeutisch wirksames nichttoxisches Säureadditionssalz oder durch Behandeln mit einer Base in ein therapeutisch wirksames nichttoxisches Basenadditionssalz umwandelt oder umgekehrt die Säureadditionssalzform durch Behandeln mit Alkali in die freie Base oder das Basenadditionssalz durch Behandeln mit Säure in die freie Säure umwandelt und gegebenenfalls deren stereochemisch isomere Formen oder  $N$ -Oxidformen herstellt.

## Revendications

## 1. Composé de formule



forme *N*-oxyde, sel d'addition pharmaceutiquement acceptable ou forme stéréochimiquement isomère de celui-ci, dans laquelle

*n* vaut 0, 1 ou 2 ;

*m* vaut 1 ou 2, à condition que si *m* vaut 2, alors *n* vaut 1 ;

*p* vaut 1 ou 2 ;

*=Q* est =O ou =NR<sup>3</sup> ;





*X* est une liaison covalente ou un radical bivalent de formule -O-, -S-, -NR<sup>3</sup>- ;

*R*<sup>1</sup> est Ar<sup>1</sup>, Ar<sup>1</sup>C<sub>1-6</sub>-alkyle ou di(Ar<sup>1</sup>)C<sub>1-6</sub>-alkyle, où chaque groupe alkyle en C<sub>1-6</sub> est éventuellement substitué par un hydroxy, un alkyloxy en C<sub>1-4</sub>, un oxo ou un substituant oxo cétalisé de formule -O-CH<sub>2</sub>-CH<sub>2</sub>-O ou -O-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O- ;


*R*<sup>2</sup> est Ar<sup>2</sup>, Ar<sup>2</sup>C<sub>1-6</sub>-alkyle, Het<sup>1</sup> ou Het<sup>1</sup>C<sub>1-6</sub>-alkyle ;

*R*<sup>3</sup> est un hydrogène ou un alkyle en C<sub>1-6</sub> ;

*L* est un hydrogène ; Ar<sup>3</sup> ; un alkyle en C<sub>1-6</sub> ; un alkyle en C<sub>1-6</sub> substitué par 1 ou 2 substituants choisis parmi un hydroxy, un alkyloxy en C<sub>1-6</sub>, Ar<sup>3</sup>, Ar<sup>3</sup>C<sub>1-6</sub>-alkyloxy et Het<sup>2</sup> ; un alcényle en C<sub>3-6</sub> ; Ar<sup>3</sup>C<sub>3-6</sub>-alcényle ; di(Ar<sup>3</sup>)C<sub>3-6</sub>-alcényle ou un radical de formule



ou



dans laquelle

- chaque q vaut indépendamment 2, 3 ou 4 ;  
 chaque r vaut 0, 1, 2, 3 ou 4 ;  
 chaque Y<sup>1</sup> est indépendamment une liaison covalente, -O- ou NR<sup>3</sup> ;  
 Y<sup>2</sup> est une liaison covalente, un C<sub>1-4</sub>-alcanediyle ou -C<sub>1-4</sub>-alkylNR<sup>3</sup>- ;  
 chaque -A=B- est indépendamment un radical bivalent de formule -CH=CH-, -N=CH- ou -CH=N- ;  
 chaque R<sup>4</sup> est indépendamment un hydrogène, un alkyle en C<sub>1-6</sub>, Ar<sup>2</sup> ou Ar<sup>2</sup>C<sub>1-6</sub>-alkyle ;  
 R<sup>5</sup> est un hydrogène, un alkyle en C<sub>1-6</sub> ou Ar<sup>3</sup> ;  
 R<sup>6</sup> est un alkyle en C<sub>1-6</sub>, Ar<sup>3</sup>, Ar<sup>3</sup>C<sub>1-6</sub>-alkyle, di(Ar<sup>3</sup>)C<sub>1-6</sub>-alkyle, Ar<sup>3</sup>C<sub>3-7</sub>-cycloalkyle, ou un indolye ;  
 R<sup>7</sup> est Ar<sup>3</sup>, Ar<sup>3</sup>C<sub>1-6</sub>-alkyle ; di(Ar<sup>3</sup>)C<sub>1-6</sub>-alkyle ; un alkyle en C<sub>1-6</sub> ; un cycloalkyle en C<sub>3-7</sub> ; un cycloalkyle en C<sub>3-7</sub> substitué par Ar<sup>3</sup> ; un oxazolyle ; un oxazolyle substitué par un halogéno ou un alkyle en C<sub>1-6</sub> ; un thiazolyle ; un thiazolyle substitué par un halogéno ou un alkyle en C<sub>1-6</sub> ; un imidazolyle ; un imidazolyle substitué par Ar<sup>3</sup>, un alkyle en C<sub>1-6</sub>, Ar<sup>3</sup>C<sub>1-6</sub>-alkyle ou un halogéno ; un indolinyle ; un indolinyle substitué par un alkyle en C<sub>1-4</sub> ; un 2,3,4-trihydroquinoléinyle ; un pyrrolidinyle ou un furanyle ;  
 chaque R<sup>8</sup> est indépendamment un hydrogène, un alkyle en C<sub>1-6</sub>, un cycloalkyle en C<sub>3-7</sub> ou un radical de formule



ou



dans laquelle Alk est un C<sub>1-6</sub>-alcanediyle ;

- Z est un radical bivalent de formule -O-, -S- ou -NR<sup>3</sup>- ;  
 R<sup>11</sup> est un phényle ; un phényle substitué par 1 ou 2 substituants choisis parmi un halogéno, un alkyle en C<sub>1-6</sub> ou un alkyloxy en C<sub>1-6</sub> ; un furanyle ; un furanyle substitué par 1 ou 2 substituants choisis parmi un alkyle en C<sub>1-6</sub> ou un hydroxy-C<sub>1-6</sub>-alkyle ; un thiényle ; un thiényle substitué par 1 ou 2 substituants choisis parmi un halogéno ou un alkyle en C<sub>1-6</sub> ; un oxazolyle ; un oxazolyle substitué par 1 ou 2 substituants alkyle en C<sub>1-6</sub> ; un thiazolyle ; un thiazolyle substitué par 1 ou 2 substituants alkyle en C<sub>1-6</sub> ; un pyridinyle ou un pyridinyle substitué par 1 ou 2 substituants alkyle en C<sub>1-6</sub> ;  
 R<sup>12</sup> est un alkyle en C<sub>1-6</sub> ou un alkyle en C<sub>1-6</sub> substitué par un hydroxy, un carboxy ou un C<sub>1-6</sub>-alkyloxy-carbonyle ;  
 Ar<sup>1</sup> est un phényle ; un phényle substitué par 1, 2 ou 3 substituants chacun choisi indépendamment parmi un halogéno, un alkyle en C<sub>1-4</sub>, un halogéno C<sub>1-4</sub>-alkyle, un cyano, un aminocarbonyle, un alkyloxy en C<sub>1-4</sub> ou un halogéno en C<sub>1-4</sub>-alkyloxy ;  
 Ar<sup>2</sup> est un naphtalényle ; un phényle ; un phényle substitué par 1, 2 ou 3 substituants chacun choisi indépendamment parmi un hydroxy, un halogéno, un cyano, un nitro, un amino, un mono- ou un di-(C<sub>1-4</sub>-alkyl) amino, un alkyle en C<sub>1-4</sub>, un halogéno C<sub>1-4</sub>-alkyle, un alkyloxy en C<sub>1-4</sub>, un halogéno C<sub>1-4</sub>-alkyloxy, un carboxy, un C<sub>1-4</sub>-alkyloxy-carbonyle, un aminocarbonyle et un mono- ou un di-(C<sub>1-4</sub>-alkyl)aminocarbonyle ;  
 Ar<sup>3</sup> est un phényle ou un phényle substitué par 1, 2 ou 3 substituants choisis parmi un halogéno, un hydroxy, un amino, un nitro, un aminocarbonyle, un alkyle en C<sub>1-6</sub>, un halogéno C<sub>1-6</sub>-alkyle ou un alkyloxy en C<sub>1-6</sub> ;  
 Het<sup>1</sup> est un hétérocycle monocyclique choisi parmi un pyrrolyle, un pyrazolyle, un imidazolyle, un furanyle, un thiényle, un oxazolyle, un isoxazolyle, un thiazolyle, un isothiazolyle, un pyridinyle, un pyrimidinyle, un



pyrazinyle et un pyridazinyle ; ou un hétérocycle bicyclique choisi parmi un quinoléinyle, un quinoxalinyne, un indolyle, un benzimidazolyle, un benzoxazolyle, un benzisoxazolyle, un benzothiazolyle, un benzisothiazolyle, un benzofuranyne et un benzothiényne ; chaque hétérocycle monocyclique et bicyclique peut éventuellement être substitué sur un atome de carbone par 1 ou 2 substituants choisis parmi un halogéno,

un alkyle en C<sub>1-4</sub> ou un mono- un di- ou un trihalogénométhyle ; et  
 Het<sup>2</sup> est un hétérocycle choisi parmi un 1,4-dihydro-5-oxo-tétrazol-1-yle, un imidazo[1,2-a]pyridinyle, un oxazolyle ou un imidazolyle ; chacun desdits hétérocycliques peut être éventuellement substitué par 1 ou, lorsque cela est possible, 2 substituants choisis parmi un alkyle en C<sub>1-4</sub> et Ar<sup>3</sup>.

2. Composé selon la revendication 1, dans lequel L est un hydrogène ; un alkyle en C<sub>1-6</sub> ; un alkyle en C<sub>1-6</sub> substitué par un hydroxy ; un alcényle en C<sub>3-6</sub> ; Ar<sup>3</sup> ; Ar<sup>3</sup>C<sub>1-6</sub>-alkyle ; di(Ar<sup>3</sup>)C<sub>1-6</sub>-alkyle ; Ar<sup>3</sup>C<sub>3-6</sub>-alcényle ; di(Ar<sup>3</sup>)C<sub>1-6</sub>-alcényle ; ou un radical de formule (a-1), (a-2), (a-4) ou (a-5) dans laquelle

R<sup>7</sup> est Ar<sup>3</sup> ; Ar<sup>3</sup>C<sub>1-6</sub>-alkyle ; di(Ar<sup>3</sup>)C<sub>1-6</sub>-alkyle ; un alkyle en C<sub>1-6</sub> ; un cycloalkyle en C<sub>3-7</sub> ; un cycloalkyle en C<sub>3-7</sub> substitué par Ar<sup>3</sup> ; un oxazolyle ; un oxazolyle substitué par un halogéno ou un alkyle en C<sub>1-6</sub> ; un thiazolyle ; un thiazolyle substitué par un halogéno ou un alkyle en C<sub>1-6</sub> ; un imidazolyle ; un imidazolyle substitué par Ar<sup>3</sup>, un alkyle en C<sub>1-6</sub>, Ar<sup>3</sup>C<sub>1-6</sub>-alkyle ou un halogéno ; un pyrrolidinyle ou un furanyne ;

Ar<sup>3</sup> est un phényle ou un phényle substitué par 1, 2 ou 3 substituants choisis parmi un halogéno, un hydroxy, un amino, un aminocarbonyl, un alkyle en C<sub>1-6</sub>, un halogénoC<sub>1-6</sub>-alkyle ou un alkyloxy en C<sub>1-6</sub> ;

Het<sup>1</sup> est un hétérocycle monocyclique choisi parmi un pyrrolyle, un pyrazolyle, un imidazolyle, un furanyne, un thiényne, un oxazolyle, un isoxazolyle, un thiazolyle, un isothiazolyle, un pyridinyle, un pyrimidinyle, un pyrazinyle et un pyridazinyle ; ou un hétérocycle bicyclique choisi parmi un quinoléinyle, un benzimidazolyle, un benzoxazolyle, un benzisoxazolyle, un benzothiazolyle, un benzisothiazolyle, un benzofuranyne et un benzothiényne ; chaque hétérocycle monocyclique et bicyclique peut éventuellement être substitué sur un atome de carbone par 1 ou 2 substituants choisis parmi un halogéno, un alkyle en C<sub>1-4</sub> ou un mono-, un di- ou un trihalogénométhyle.

3. Composé selon la revendication 1 ou 2, dans lequel R<sup>1</sup> est Ar<sup>1</sup>C<sub>1-6</sub>-alkyle, R<sup>2</sup> est un phényle substitué par 2 substituants choisis parmi un méthyle et un trifluorométhyle, X est une liaison covalente et =Q est =O.

4. Composé selon l'une quelconque des revendications 1 à 3, dans lequel n et m valent 1 et p vaut 1 ou 2.

5. Composé selon l'une quelconque des revendications 1 à 4, dans lequel R<sup>1</sup> est un phénylméthyle ; R<sup>2</sup> est un phényle substitué par 2 substituants choisis parmi un méthyle ou un trifluorométhyle ; n, m et p valent 1 ; X est une liaison covalente ; et =Q est =O.

6. Composé selon l'une quelconque des revendications 1 à 4, dans lequel L est un radical de formule (a-2) dans laquelle R<sup>4</sup> est un hydrogène ou un phényle ; r vaut 0 ou 1 ; Y<sup>1</sup> est une liaison covalente, -O- ou -NH-, R<sup>7</sup> est un pyrrolidinyle ; un furanyne ; un 1-phénylcyclohexanyne ; un diphénylméthyle ; ou un phényle substitué par 1, 2 ou 3 substituants choisis chacun indépendamment parmi un méthyle, un méthoxy ou un chloro.

7. Composé selon la revendication 5 ou 6, dans lequel le composé est de configuration trans.

8. Composé selon la revendication 5 ou 6, dans lequel le composé est de configuration cis.

9. Composé selon la revendication 1, dans lequel L est un hydrogène.

10. Composé selon la revendication 1, le composé étant

le 4-[1-[3,5-bis(trifluorométhyle)benzoyl]-2-(phénylméthyle)-4-pipéridinyle]-N-(2,6-diméthylphényl)-1-pipérazine-acétamide ;

le 4-[1-[3,5-bis(trifluorométhyle)benzoyl]-2-(phénylméthyle)-4-pipéridinyle]-N-(1-phénylcyclohexyle)-1-pipérazine-acétamide ;

le 1-[3,5-bis(trifluorométhyle)benzoyl]-2-(phénylméthyle)-4-[4-[α-(1-pyrrolidinylcarbonyl)benzyle]-1-pipérazinyle]pipéridine ;

le 1-[3,5-bis(trifluorométhyle)benzoyl]-4-[4-[1-[(2-méthyle-5-oxazolyle)méthyle]-1H-benzimidazol-2-yl]-1-pipérazinyle]-2-(phénylméthyle)pipéridine ;

le 4-[1-[3,5-bis(trifluorométhyl)benzoyl]-2-[(4-trifluorométhylphényl)méthyl]-4-pipéridinyl]-N-(2,6-diméthylphényl)-1-pipérazine acétamide ;  
 le 4-[1-[3,5-bis(trifluorométhyl)benzoyl]-2-[(3,4-dichlorophényl)méthyl]-4-pipéridinyl]-N-(2,6-diméthylphényl)-1-pipérazine acétamide.

11. Composé selon la revendication 10, le composé

étant le (+)-(B)-*trans*-4-[1-[3,5-bis(trifluorométhyl)benzoyl]-2-(phénylméthyl)-4-pipéridinyl]-N-(2,6-diméthylphényl)-1-pipérazine acétamide ;  
 le (-)-(B)-*cis*-4-[1-[3,5-bis(trifluorométhyl)benzoyl]-2-(phénylméthyl)-4-pipéridinyl]-N-(2,6-diméthylphényl)-1-pipérazine acétamide ; ou  
 le (+)-(B)-*trans*-4-[1-[3,5-bis(trifluorométhyl)benzoyl]-2-(phénylméthyl)-4-pipéridinyl]-N-(2,6-diméthylphényl)-1-pipérazine acétamide, l'acide(L)-malique (1:1).


12. Composition comprenant un support pharmaceutiquement acceptable et, à titre d'ingrédient actif, une quantité thérapeutiquement efficace d'un composé selon l'une quelconque des revendications 1 à 11.

13. Procédé de préparation d'une composition selon la revendication 12, caractérisé en ce qu'un support pharmaceutiquement acceptable est mélangé intimement avec une quantité thérapeutiquement efficace d'un composé selon l'une quelconque des revendications 1 à 11.

14. Composé selon l'une quelconque des revendications 1 à 11, destiné à être utilisé en tant que médicament.


15. Procédé de préparation d'un composé selon la revendication 1, caractérisé par

a) la N-alkylation réductive d'un intermédiaire de formule (II) dans laquelle L et p sont tels que définis à la revendication 1, avec un intermédiaire de formule (I)




dans laquelle R<sup>1</sup>, R<sup>2</sup>, X, Q, n et m sont tels que définis à la revendication 1, dans un solvant inerte vis-à-vis de la réaction, en présence d'un réducteur approprié et éventuellement en présence d'un agent complexant approprié ;

b) la réaction d'un intermédiaire de formule (IV) dans laquelle R<sup>2</sup>, X et Q sont tels que définis à la revendication 1 et W<sup>1</sup> est un groupe partant approprié, avec un intermédiaire de formule (V)




dans laquelle R<sup>1</sup>, L, n, m et p sont tels que définis à la revendication 1, dans un solvant inerte vis-à-vis de la réaction et en présence d'une base appropriée ;

c) la N-alkylation réductive d'un dérivé pipérazinique de formule (VII) dans laquelle p est tel que défini à la revendication 1 et P<sup>1</sup> est un groupe protecteur, avec un intermédiaire de formule (II)




dans laquelle  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{X}$ ,  $\text{Q}$ ,  $n$  et  $m$  sont tels que définis à la revendication 1, dans un solvant inerte vis-à-vis de la réaction, en présence d'un réducteur approprié et éventuellement en présence d'un agent complexant approprié, en formant ainsi un composé de formule (I-c) ;

d) la *N*-alkylation réductive d'un dérivé pipérazinique de formule (VII) dans laquelle  $p$  est tel que défini à la revendication 1 et  $\text{P}^1$  est un groupe protecteur, avec un intermédiaire de formule (VIII),




dans laquelle  $\text{R}^1$ ,  $n$  et  $m$  sont tels que définis à la revendication 1, dans un solvant inerte vis-à-vis de la réaction et en présence d'une base appropriée, en formant ainsi un intermédiaire de formule (XI) ; qui peut ensuite être mis à réagir avec un intermédiaire de formule (IV)




dans laquelle  $\text{W}^1$  est un groupe partant approprié et  $\text{X}$ ,  $\text{Q}$  et  $\text{R}^2$  sont tels que définis à la revendication 1, pour former un composé de formule (I-c) ;

e) la déprotection d'un composé de formule (I-c) en employant des techniques de déprotection connues dans l'art, en formant ainsi un composé de formule (I-b) ;



f) la réaction d'un composé de formule (I-b) avec un intermédiaire de formule (VI)



dans laquelle L' est identique à L tel que défini à la revendication 1 mais est différent d'un hydrogène, et W<sup>2</sup> est un groupe partant approprié, dans un solvant inerte vis-à-vis de la réaction et en présence d'une base appropriée ; en formant ainsi un composé de formule (I-a) ;

et, si on le souhaite, la transformation des composés de formule (I) les uns en les autres en suivant des transformations connues dans l'art et en outre, si on le souhaite, la transformation des composés de formule (I) en un sel d'addition à un acide thérapeutiquement actif et non toxique par traitement avec un acide, ou en un sel d'addition à une base thérapeutiquement actif et non toxique par traitement avec une base, ou inversement, la transformation de la forme de sel d'addition à un acide en base libre par traitement avec un alcali, ou la transformation du sel d'addition à une base en acide libre par traitement avec un acide ; et si on le souhaite, la préparation de formes stéréochimiquement isomères de formes N-oxyde de ceux-ci.